

FLORA AND FAUNA
WITHIN THIRTY MILES OF
HANOVER, N. H.

QK
112
J 47



Republican Press Ass'n,
Printing and Binding,
Concord, N. H.



MBL/WHOI



0 0301 0013504 2

A CATALOGUE
OF THE
FLOWERING PLANTS
AND
HIGHER CRYPTOGAMS,

BOTH NATIVE AND INTRODUCED, FOUND WITHIN ABOUT THIRTY
MILES OF HANOVER, N. H., INCLUDING A FEW CULTI-
VATED SPECIES, TO WHICH IS APPENDED

A List of Vertebrate Animals of the
Same Region.

BY
HENRY G. JESUP, A. M.,
PROFESSOR OF NATURAL HISTORY.

PRINTED FOR THE AUTHOR.

1891.

Directions for Collecting and Preserving Plants and for the Formation of an Herbarium.

1. A *complete* specimen will represent the root, stem, leaves, flowers, and fruit.
 2. Collect small plants, like the Violet, *entire*.
 3. When the leaves of the stem and the root differ, as in some Buttercups, get both.
 4. If the staminate and the pistillate flowers are on different plants, as in Thalictrum (Rue), or on different parts of the same plant, as in the Butternut, look for both.
 5. Plants of the Mustard Family (Cruciferæ) and the Parsley Family (Umbelliferæ), with all Rushes and Sedges, are classified by the *fruit*. Grasses may be gathered in *flower*.
 6. Ferns should be in fruit, and when the sterile and fertile fronds differ, as in Osmunda, get both.
 7. Thick roots and stems can be split in two, and one half only be preserved.
 8. Arrange the plant to be dried in a folded sheet of thin, bibulous paper, and do not disturb it until dry. Let thick pads of drying paper alternate with these single sheets, and then place the whole under pressure. Change the drying pads every day or two, and dry as rapidly as practicable.
 9. These dried plants attached to sheets of paper by strips of surgeon's plaster or by glue, fully ticketed with name, when and where and by whom collected, and arranged in Orders and Genera as in the Manual, constitute an Herbarium.
-

Summary of Species.

Polypetals	323
Gamopetals	312
Apetals	111
Angiosperms	—746
Gymnosperms	15
Exogens	—761
Endogens	335
Phænogamis	—1096
Cryptogams	65
Total number of species in the Flora proper	—1161
Cultivated species and genera	115

1203

P R E F A C E.

The following Catalogue is intended to include, so far as known, the Flora within about thirty miles radius of the town of Hanover, New Hampshire, a circuit extending to and embracing the mountain peaks Moosilauke and Kearsarge in New Hampshire, and Killington in Vermont. The outlying towns intersected by the circumference of the circle, beginning with the southern limit and the Connecticut river, are, in New Hampshire, Charlestown, Acworth, Lempster, Goshen, Newbury, Sutton, Andover, Hill, New Hampton, Bristol, Bridgewater, Plymouth, Campton, Ellsworth, Benton, and Haverhill. Continuing the circuit on the Vermont side of the river, the towns are Newbury, Topsham, Orange, Williamstown, Brookfield, Braintree, Rochester, Pittsfield, Chittenden, Mendon, Shrewsbury, Mt. Holly, Ludlow, Chester, and Springfield. The region thus indicated is divided nearly equally between the two adjoining states by their common boundary the Connecticut river, and includes a variety of soil, climate, and location sufficient to furnish a home for a majority of the plants of northern New England, except those peculiar to the sea-shore and the higher mountains.

The only river of any considerable size is the Connecticut, running from north to south throughout the entire district, a distance of sixty miles or more. Numerous tributaries join this river on both sides, the largest of which, in Vermont, are the Ompompanoosuc which empties in Norwich, the White river at White River Junction, the Ottoquechee in Hartland, a few miles from whose mouth is the deep gorge known as "Quechee Gulf," and the Black river in Springfield. In New Hampshire, the Mascoma, the outlet of Mascoma Lake, joins the Connecticut in West Lebanon, and the Sugar river in Claremont. The Pemigewasset and other streams which form the head waters of the Merrimack river, moreover, reach some towns in the eastern part of the district.

The larger sheets of water included are, in New Hampshire, Sunapee lake, Newfound lake, and Mascoma lake, or Enfield pond. Sunapee lake, in the towns of Sunapee, Newbury, and New London, is about seven miles long, and at one point two and one half miles broad, and is a well known summer resort. Newfound lake in the towns of Hebron, Bridgewater, and Bristol, is less known, but is a fine sheet of water, six miles perhaps in length, and three miles in extreme breadth, and very nearly as large as Sunapee. Mascoma lake or Enfield pond is somewhat smaller, being four and a half miles long by rather more than half a mile in width.

On the Vermont side there is little to correspond with the above. Fairlee pond or Morey lake in the town of Fairlee, and Fairlee lake, partly in the same town, are both comparatively small. The purity of the water in these larger lakes, and their commonly bold or gravelly shores, are not favorable to the accumulation of much aquatic vegetation, but even here the botanist will

find some plants, like *Bidens Beckii* (Water Marigold) for instance, not found elsewhere, and all these lakes will repay exploration. It is the smaller ponds and adjoining marshes, rich with decaying vegetation, that more abundantly reward the persevering herbalist. Of these, also, there are more on the eastern side of the Connecticut river than on the western side. Not a few of them are high up among the hills, visited very seldom and by very few, but in these high and cool bogs there still is found the remains of a vegetation that long since deserted the warmer lowlands. Here are the rarer Sedges and Grasses, *Sarracenia* (Pitcher Plant), and Orchids, besides Ericaceous plants like *Azalea*, *Rhodora*, *Andromeda*, and *Ledum* (Labrador Tea), as well as others of equal interest. Periods of drought seldom reach these rich and moist garden spots.

Of the three mountain peaks referred to above, Moosilauke in Benton, N. H., 4,811 ft. in altitude, is the only one having a treeless summit of any considerable extent, suggestive of Alpine plants, and there a few species appear, such as *Arenaria Groenlandica* (Mountain Sandwort), *Loiseleuria procumbens* (Alpine Azalea), *Vaccinium Vitis-Idaea* (Mountain Cranberry), *Vaccinium uliginosum* (Bog Bilberry), and *Solidago alpina* (Alpine Golden-rod), etc. Kearsarge Mt. near Salisbury, N. H., and Killington Mt., with the adjoining peaks in Sherburne and Mendon, Vt., are well wooded nearly to their summits, and have upon them little or nothing strictly alpine. The same is true of Mt. Ascutney near Windsor, Vt., and the same may be said of Cube Mt. in Orford, and Smart's Mt. in Lyme, N. H. This latter elevation is peculiar in this, that the timber on the summit, instead of being stunted and worthless, is of full marketable size. Cardigan Mt. in Orange, N. H., has a conspicuous treeless summit, but little truly Alpine, unless we except *Vaccinium Vitis-Idaea*, has been reported from there. It is probable that a more careful exploration might result in some new discoveries on any or all of these mountains.

In addition to the lakes and mountains, numerous streams are found within the district on whose banks and in whose waters are forms of vegetation peculiar to such localities and to them alone. Rivers like the Connecticut, with numerous tributaries, are great highways for the distribution of both seeds and growing plants. Every freshet bears these along in its swift current, and transports them many miles from the place of their origin. They are stranded on the shore, or find a home in the still pools along the margin of the stream, and once established may remain indefinitely. This explains the presence of *Astragalus alpinus* (Alpine-Vetch) on the rocks at Sumner's Falls in Plainfield, N. H.—a plant of the far north—together with *Astragalus Robbinsii*, a species of equal interest. When, moreover, the river recedes during a period of drought, the shore that is laid bare supplies conditions under which a still different vegetation makes its appearance. *Ranunculus Flammula, var. reptans* (Creeping Buttercup) then spreads its green carpet dotted with yellow stars upon the sand; and other plants, that are shut out of their needed supply of oxygen by high water, begin once more to flourish.

Plants are as migratory as any other of the world's inhabitants. They are always ready to make use of the same great thoroughfares and of the same

means of transportation. Every new railroad or canal or highway opens for them a direct line of communication with regions not yet visited, and they instantly make use of them all. Every steamship or sailing vessel, whether from across the ocean or coasting along our own shores, aids in the distribution, while all these and similar agencies are but supplementary to those which, entirely apart from civilization, are in constant operation.

The early inhabitants of New England had scarcely landed on our shores before they found that the Ox-eye Daisy and the Dandelion, together with the Buttercups and Docks and Plantains and a score of similar emigrants from the old world, had accompanied them. And when the tide of travel advanced westward toward the Mississippi river, not only did these same plants follow the same route, but in due time the plants of the prairie, like *Rudbeckia hirta* (Yellow Daisy or Cone-flower) began to appear in Eastern meadows. These introduced plants are sometimes of great value, like the Grasses and Clovers, but commonly they are the so called weeds of the farm and the garden, and come unbidden and unwelcome. Some come to stay, others that appear occasionally about our manufacturing establishments, persist for a time only, but finally disappear to be followed very likely by a new importation. Plants of this kind are in some respects more interesting than those of our native Flora, not only because we come more directly in contact with them, but because each has a history behind it and around it, which serves to illustrate the beginnings and the development of that portion of the country where it has found a new home.

The fields and open woodlands, the roadsides and fence rows, are naturally first visited by the plant collector because easily accessible, but every nook and corner, whether in sunshine or in shade, in its season, has its own share of vegetable as well as animal life, and the lover of Nature who would become thoroughly acquainted with her treasures seeks them everywhere, for the simple reason that they may be found everywhere. The Flora of no section of our country will ever be known so completely that nothing more shall remain to be learned about it. No surprising discoveries of new species may be made, though of this we dare not be too sure, but new chapters in the history of well known plants may be opened, newly arrived immigrants noted, new varieties found, new facts as to distribution and new economical uses ascertained, as well as new relations between the species and its environment in illustration of the conflict all plants are making successfully or unsuccessfully for an existence. For reasons like these a region which has been as long settled and is as well known as the one embraced in this Catalogue will never be without interest to the naturalist.

The relation of what is here called the Hanover Flora to the general New England Flora calls for some brief notice. Thirty miles south of Hanover, in the towns of Charlestown, N. H., and Springfield, Vt., some of the trees and shrubs so common a little farther south in both of these states have already reached their northern limit, and begin to disappear from the native woods. The Chestnut and the Shagbark Hickory are no longer abundant. The season is too short for any considerable quantity of fruit to reach perfection, and seedlings are rare. Trees of these two kinds when planted in sheltered localities

may attain considerable age and size. A single chestnut of exceptional size may be seen in Norwich, Vt., five feet eight inches in diameter, and more than fifty years old, but it was planted where it now stands, and the chestnuts are of little value. An occasional Shagbark Hickory of moderate size may also be met with, though the Pignut Hickory is not infrequent in the woods, and the Bitternut Hickory ascends the Connecticut as far north as Wells river. The Black Birch, and the Grey Birch, as well as the Pitch Pine, are rare north of Hanover and not common in its vicinity, while the Sycamore or Buttonwood is nowhere seen in a wild state. To the above may be added as decidedly rare, the Sassafras, the Spice Bush (*Lindera*), the Flowering Dogwood (*Corneus florida*), the Red Cedar, and the common Juniper, together with many other trees and shrubs that fifty miles south of Hanover may be common.

On the other hand, as the traveller advances thirty miles northward from Hanover, another class of aborescent vegetation appears, that either is not found at all southward, or at least is not abundant. Here the White Cedar becomes plentiful and very valuable, the Tamarack or American Larch fills the swamps, and the White Spruce occasionally appears, while the Black Spruce covers the lower mountain tops, the Paper Birch adorns the hillsides, the Green Alder is found by the streams and in the ravines, and the Elder has red berries more commonly than black ones. Additional illustrations might readily be given, but the above will indicate what changes may be noted within sixty miles only of latitude.

The first and preliminary edition of this Catalogue was issued in 1882. Since that time some additional explorations of the region have been made, very much increasing the list of names and supplying much valuable information.

In order to make the Catalogue as useful as possible to the author's classes in Natural History, for whom primarily it has been prepared, there have been added the names of such plants as are most commonly met with in cultivation, arranged under their respective Orders. These do not properly belong to the Flora, and are distinguished from the rest either by the arrangement of the type or by being enclosed in square brackets. Their description will be found in *Gray's Field, Forest, and Garden Botany*, while the description of plants belonging to the Flora proper will be found in *Gray's Manual of the Botany of the Northern United States, Sixth Edition, 1890*. The synonyms refer to the fifth edition of the same Manual. The names of introduced and cultivated species are printed in SMALL CAPITALS. The local or English name of each plant has been added so far as known. And as an additional aid to the student, brief directions have been given for collecting and preserving plants and for the formation of an Herbarium. Localities not designated are in New Hampshire.

In the preparation of the previous edition of the Flora, free use was made of a Catalogue of New Hampshire plants compiled for the Geological Survey of the State by Wm. F. Flint, and a large amount of information and personal assistance was contributed by Nathan Barrows, M. D., then of Meriden. From the Vermont side of the Connecticut river some rare species were

reported from North Pomfret by Prof. A. P. Morgan; from Sharon by E. K. Baxter, M. D.; and from Union Village by Ferdinand Blanchard, M. D.; while Mt. Ascutney and the vicinity of Windsor were thoroughly explored by George H. Leland.

For the present edition it has been found necessary to rewrite and rearrange a large part of the Flora to correspond with the changes in both classification and nomenclature found in the last edition of the Manual referred to above. As the work has been in progress, important aid has generously been furnished by the author's numerous friends and correspondents in various parts of the country—too numerous to be mentioned individually—and he would here convey to them his hearty thanks. The students who year by year have been connected with his classes have done their share also in bringing this Flora to its present completeness. The names of very many of these contributors, both correspondents and students, will be found in their appropriate places in the following pages.

The List of Vertebrate Animals found within the same district as the Flora, originally given in the edition of 1882, was prepared by Prof. T. W. D. Worthen, with the assistance of Prof. B. T. Blanpied and Prof. C. S. Cook, and was intended for use with Jordan's Vertebrates of the Northern United States as published in 1880.

The List here given has also been prepared by Prof. Worthen who has embodied his own notes made since the previous edition was issued, as well as some additional information furnished by other naturalists, residents now or of late in this vicinity, whose names are given in connection with the items reported by them. To Prof. Worthen and to these gentlemen cordial thanks are rendered.

The classification adopted is that of the *Fifth and Revised Edition of Jordan's Vertebrates of 1888*, involving very many changes in both names and general arrangement. The above book must therefore be consulted for full descriptions.

The accompanying map, contributed by the students, indicates clearly the region represented by the Flora and Fauna.

Additions and corrections are solicited, and will always be welcome.

H. G. J.

Hanover, N. H., Jan. 23, 1891.



MAP OF THE REGION WITHIN THIRTY MILES OF HANOVER, N. H.



CATALOGUE OF PLANTS.

SERIES 1.

PHÆNOGAMOUS OR FLOWERING PLANTS.

CLASS I. DICOTYLEDONOUS OR EXOGENOUS PLANTS.

SUBCLASS I. ANGIOSPERMÆ.

DIVISION I. POLYPETALOUS EXOGENS.

ORDER I. RANUNCULACEÆ. CROWFOOT FAMILY.

Clematis, L. Virgin's Bower.

C. Virginiana, L. River-banks, etc.; common.

C. verticillaris, DC. Open woods. Windsor, Vt., G. H. Leland; Hartland, Vt., W. W. Eggleston; Lyme, N. H., Z. A. Norris; rare.

Anemone, Tourn. Wind-Flower. Anemone.

A. cylindrica, Gray. Pomfret, Windsor, and Norwich, Vt.; not abundant.

A. Virginiana, L. Woods and roadsides; common.

A. nemorosa, L. Margin of woods; common.

Hepatica, Dill. Liver-leaf. Liverwort.

H. triloba, Chaix. Woods. More common southward.

H. acutiloba, DC. Woods. The more abundant species.

Anemonella, Spach. Rue-Anemone.

A. thalictroides, Spach. (*Thalictrum anemonoides*, Michx.) Claremont, Dr. N. Barrows; rare.

Thalictrum, Tourn. Meadow-Rue.

T. dioicum, L. (Early-Rue.) Rocky woods; common.

T. polygamum, Muhl. (T. Cornuti, Gray's Manual, 5th ed.)—(Late-Rue.) Wet meadows; common.

Ranunculus, Tourn. Crowfoot. Buttercup.

- R. circinatus*, Sibth. (*R. divaricatus*, Manual, 5th ed.)—(White Water-Crowfoot.) Cold streams and ponds. Windsor, Vt., *Leland*; rare.
R. aquatilis, L., var. *trichophyllum*, Gray. (White Water-Crowfoot.) Slow streams. *Dr. Barrows*; rare.
R. multifidus, Pursh. (Yellow Water-Crowfoot.) Hanover, N. H.; Norwich, Vt.; rare.
R. Flammula, L., var. *reptans*, E. Meyer. (Small Creeping-Crowfoot) Sandy shore of Conn. river; shore of pond, New London, *H. E. Sargent*.
R. abortivus, L. (Small-flowered Crowfoot.) Common.
R. sceleratus, L. North Charlestown, *Dr. Graves*; rare.
R. recurvatus, Poir. (Hooked Crowfoot.) Common.
R. fascicularis, Muhl. (Early C.) North Charlestown, *Dr. Graves*.
R. septentrionalis, Poir. (*R. repens*, Manual, 5th ed.)—(Large Creeping-Crowfoot.) Wet meadows; common.
R. Pennsylvanicus, L. (Bristly Crowfoot.) Wet places; common.
R. BULBOSUS, L. Hanover, *F. P. Comings*; Meriden, *Dr. Barrows*; Norwich, Vt., *Dr. Blanchard*; Chelsea, Vt., *J. M. Comstock*.
R. ACRIS, L. Common Buttercup. This and the above, European weeds.

Caltha, L. Marsh-Marigold. Cowslips.

- C. palustris*, L. Hanover and Lebanon, N. H.; Norwich, Vt.; etc.; common.

Coptis, Salisb. Gold-thread.

- C. trifolia*, Salisb. Wet woods; common.

Aquilegia, Tourn. Columbine.

- A. Canadensis*, L. Rocks; common.

Cimicifuga, L. Bugbane.

- C. racemosa*, Nutt. (Black Cohosh.) Claremont, *Dr. Barrows*; very rare.

Actaea, L. Baneberry.

- A. spicata*, L., var. *rubra*, Ait. (Red Baneberry.) Rich woods; common.
A. alba, Bigel. (White Baneberry.) A little later in flowering.

[Among cultivated plants of this Order are various species of ANEMONE, CLEMATIS, AQUILEGIA (Columbine), DELPHINIUM (Larkspur), ACONITUM (Monkshood), PÆONIA (Pæony), RANUNCULUS, etc.]

ORDER 2. **MENISPERMACEÆ**. MOONSEED FAMILY.**Menispermum**, L. Moonseed.

- M. Canadense*, L. Claremont, both sides Conn. river. *Dr. Barrows*.

ORDER 3. **BERBERIDACEÆ**. BARBERRY FAMILY.**Berberis**, L. Barberry.

- B. VULGARIS*, L. More commonly cultivated than wild; from Europe.

Caulophyllum, Michx. Blue Cohosh.

- C. thalictroides*, Michx. Rich woods; common.

***Podophyllum*, L.** Mandrake.

P. peltatum, L. Hanover, N. H.; Norwich, Vt., E. A. Mack : doubtless introduced.

ORDER 4. **NYMPHÆACEÆ.** WATER-LILY FAMILY.***Brasenia*, Schreber.** Water-shield.

B. peltata, Pursh. In ponds.

***Nymphaea*, Tourn.** White Water-Lily.

N. odorata, Ait. In ponds ; common.

***Nuphar*, Smith.** Yellow Pond-Lily.

N. advena, Ait. (Cow Lily.) Everywhere common.

var. minus, Morong. (N. rubrodiscum, Morong.) Squam Lake, *Jesup*. A hybrid, probably, between N. advena and the next species ; intermediate in size, with stigmas crenately toothed and bright red or crimson.

N. Kalmianum, Ait. (N. luteum, var. pumilum. Man., 5th ed.) Hanover, N. H.; Dr. Barrows ; Sherburne, Vt., W. W. Eggleston.

ORDER 5. **SARRACENIACEÆ.** PITCHER-PLANT FAMILY.***Sarracenia*, Tourn.** Pitcher-Plant. Side-Saddle Flower.

S. purpurea, L. Lebanon, Meriden, and New London, N. H.; Norwich and Sharon, Vt. In peat bogs ; not rare.

ORDER 6. **PAPAVERACEÆ.** POPPY FAMILY.***Sanguinaria*, Dill.** Blood-root.

S. Canadensis, L. Borders of rich woods ; common.

***Chelidonium*, L.** Celandine.

C. MAJUS, L. Near dwellings and fences ; from Europe.

ORDER 7. **FUMARIACEÆ.** FUMITORY FAMILY.***Adlumia*, Raf.** Mountain Fringe. Climbing Fumitory.

A. cirrhosa, Raf. Claremont and Plainfield, N. H.; Ludlow, Windsor, and Royalton, Vt.; not common ; often cultivated.

***Dicentra*, Borkh.** Dicentra.

D. Cucullaria, DC. (Dutchman's Breeches.) Rich woods ; common.

D. Canadensis, DC. (Squirrel-Corn.) Rich woods ; not rare.

[D. EXIMIA, DC., from Va. is frequent in old gardens.]

***Corydalis*, Vent.**

C. glauca, Pursh. (Pale Corydalis.) Rocky places and clearings.

C. aurea, Willd. (Golden Corydalis.) Norwich, Vt., near the Passumpsic R. R. *Jesup*.

***Fumaria*, Tourn.** Fumitory.

F. OFFICINALIS, L. A weed in gardens ; European.

ORDER 8. CRUCIFERÆ. MUSTARD FAMILY.

Dentaria, Tourn. Pepper-root.

- D. diphylla, L. Rich woods; common.
- D. laciniata, Muhl. Plainfield, *Dr. Barrows*; rare.
- D. maxima, Nutt. With the first, but less common.

Cardamine, Tourn. Bitter Cress.

- C. hirsuta, L. Wet ground. A slender variety is found in dry places.

Arabis, L. Rock-Cress.

- A. hirsuta, Scop. Rocky places; not rare.
- A. laevigata, Poir. Woods and ravines; not common.
- A. perfoliata, Lam. (Tower Mustard.) Hartford, Vt., *Flint*: Norwich, Vt., *Jesup*: Hanover, Lyme, etc.; not common.
- A. confinis, Watson. (A. Drummondii, Manual, 5th ed.) Hanover, N. H., A. C. French; Fairlee, Vt., F. O. Grover; not common.

Camelina, Crantz. False Flax.

- C. SATIVA, Crantz. Occasionally in grass grounds. European.

Nasturtium, R. Br. Water-Cress.

- N. SYLVESTRE, R. Br. (Yellow Cress.) A weed in river meadows. Norwich, Vt.; etc.; from Europe.
- N. palustre, DC. Marsh Cress. Wet places; common.
- N. ARMORACIA, Fries. (Horseradish.) Rarely seen in fruit; European.

Barbara, R. Br. Winter Cress. Yellow Rocket.

- B. vulgaris, R. Br. Low grounds; common.

Hesperis, Tourn. Common Rocket.

- H. MATRONALIS, L. Roadsides, etc.; a garden escape.

Erysimum, Tourn. Treacle Mustard.

- E. cheiranthoides, L. Hanover; a weed in gardens.

Sisymbrium, Tourn. Hedge Mustard.

- S. OFFICINALE, Scop. Common European weed.

Brassica, Tourn. Cabbage, Mustard, etc.

- B. CAMPESTRIS, L. (Kale.) Common in grain fields; a troublesome weed.
- B. SINAPISTRUM, Boissier. (English Charlock.) Plainfield, *Miss Stevens*; Plymouth, *H. E. Sargent*; not common.
- B. ALBA, Gray. (White Mustard.) Rare.
- B. NIGRA, Koch. (Black Mustard.) Common near dwellings; all from Europe.

Capsella, Medic. Shepherd's Purse.

- C. BURSA-PASTORIS, Moench. A weed in gardens; from Europe.

Thlaspi, Tourn. Pennycress.

- T. ARVENSE, L. Pomfret, Vt., *A. P. Morgan*; Lebanon, *Jesup*: waste places; rare; European.

Lepidium, Tourn. Peppergrass.

L. Virginicum, L. Roadsides; common.

L. CAMPESTRE, Br. Hanover, C. H. Dana: not common; European.

L. RUDERALE, L. Hanover, etc.; along with L. Virginicum; from Europe.

Raphanus, Tourn. Wild Radish.

R. RAPHANISTRUM, L. (Jointed Charlock.) Hanover; not rare in grain fields; European.

[Cultivated plants:—Various kinds of BRASSICA (Cabbage and Turnip), RAPHANUS SATIVUM (Garden Radish), LEPIDIUM SATIVUM (Garden Peppergrass), MATTHIOLA (Stock), IBERIS (Candytuft), ALYSSUM MARITIMUM (Sweet Alyssum), etc.]

ORDER 9. **CISTACEÆ**. ROCK-ROSE FAMILY.**Helianthemum**, Tourn. Rock-Rose.

H. Canadense, Michx. Frost-weed. Gravelly soil.

Lechea, Kalm. Pinweed.

L. major, Michx. Dry and sterile fields; common.

L. minor, L. With the above and quite variable, including perhaps more than one species.

ORDER 10. **VIOLACEÆ**. VIOLET FAMILY.**Viola**, Tourn. Violet.

V. palmata, L. (V. cucullata, var. palmata. Gray.)—(Palmately-leaved Violet.) On drier ground than the variety, and less common.

var. cucullata, Gray. (V. cucullata, Ait.)—(Common Blue Violet.) Very common everywhere.

V. sagittata, Ait. Hanover, etc. Roadsides and upland pastures; not abundant.

V. Selkirkii, Pursh. (Selkirk Violet.) Shady hillsides; not very common.

V. blanda, Willd. (Sweet White Violet.) Moist places; common, and very variable.

var. palustriformis, Gray. “The larger form; in shady, mossy ground.”

var. renifolia, Gray. (Kidney-leaved Violet.) Named from specimens collected in 1870 by F. A. Sherman: common in rich, rocky woods.

V. primulæfolia, L. (Primrose-leaved Violet.) With the next. New London, W. F. Carter: not common.

V. lanceolata, L. (Lance-leaved Violet.) Near Dorchester Line Pond, A. C. French; New London, in wet pastures, Carter: rare.

V. rotundifolia, Michx. (Round-leaved Violet.) Moist woods. The earliest species; common.

V. pubescens, Ait. (Downy Yellow Violet.) Woods; common and variable.

V. Canadensis, L. (Canada Violet.) Rich woods; common.

V. rostrata, Pursh. (Long-spurred Violet.) With the above.

V. canina, L. var. Muhlenbergii, Gray. (Dog Violet.) Everywhere common and rather variable.

V. TRICOLOR, L. (Pansy.) Spontaneous in gardens along with V. ODORATA, the English Sweet Violet.

ORDER II. CARYOPHYLLACEÆ. PINK FAMILY.

Dianthus, L. Pink.

D. ARMERIA, L. (Deptford Pink.) Randolph, Vt., *Bates*.

D. DELTOIDES, L. (Maiden Pink.) Near Sharon, Vt.; well established on the roadside, *Jesup*.

Saponaria, L. Soapwort. Bouncing Bet.

S. OFFICINALIS, L. Roadsides; commonly double; European.

Silene, L. Catchfly.

S. CUCUBALUS, Wibel. (S. inflata, Smith.)—(Bladder Campion.) Meriden and Hanover, N. H.; Chelsea, Vt.; etc. Roadsides; European.

S. antirrhina, L. (Sleepy Catchfly.) Waste places.

S. NOCTIFLORA, L. Frequent; introduced with grass seed.

[S. ARMERIA, L. (Sweet-William Catchfly.) Spontaneous in gardens.]

Lychnis, Tourn. Lychnis.

L. GITHAGO, Lam. (Corn-cockle.) Hanover, in cultivated fields; from Europe.

Arenaria, L. Sandwort.

A. SERPYLLIFOLIA, L. Windsor, Vt., *Leland*. Waste places; European.

A. Michauxii, Hook. (A. stricta, Michx.) Norwich, Vt., *Dr. Blanchard*; Fairlee Cliffs, Vt., *F. O. Grover*. Not very common.

A. Grænlandica, Spreng. (Mountain Sandwort.) Moosilauke Mt., *W. H. Flint*; Kearsarge Mt., *Dr. Barrows*.

A. lateriflora, L. Lebanon, at Olcott Falls; infrequent.

Stellaria, L. Chickweed.

S. MEDIA, Smith. (Common Chickweed.) From Europe.

S. longifolia, Muhl. (Long-leaved Stitchwort.) Grassy places.

S. longipes, Goldie. Claremont, *Dr. Barrows*; rare.

S. GRAMINEA, L. Occasional in cultivated grounds. European.

S. borealis, Bigel. Wet, shady places; not rare.

S. uliginosa, Murr. (Swamp Chickweed.) Rochester, Vt., *Eggleston*.

Cerastium, L. Chickweed.

C. VULGATUM, L. (C. viscosum of Gray's Man., 5th ed.) Common European weed.

C. arvense, L. Rocky pastures north of Windsor, Vt., *Flint*; Hanover, N. H., *G. M. Moore*; scarce.

Sagina, L. Pearlwort.

S. procumbens, L. Moist places.

Buda, Adans. Sand-Spurrey.

B. rubra, Dumort. (Spergularia rubra. Presl.) Dry, sandy soil; Canaan and Claremont, N. H., *Dr. Barrows*; Royalton, Vt., *H. Q. Ward*; Plymouth and New London, N. H., *Sargent*.

Spergula, L. Spurrey.

S. ARVENSIS, L. (Corn-Spurrey.) Hanover, Meriden, etc.; in cultivated grounds; European.

ORDER 12. **PORTULACACEÆ**. PURSLANE FAMILY.**Portulaca**, Tourn. Purslane.

P. OLERACEA, L. A common weed in gardens; European.

Claytonia, Gronov. Spring-Beauty.

C. CAROLINIANA, Michx. Common in open woods.

ORDER 13. **HYPERICACEÆ**. ST. JOHN'S-WORT FAMILY.**Hypericum**, Tourn. St. John's-wort.

H. ASCYRON, L. (*H. PYRAMIDATUM*, Ait.)—(Great St. John's-wort.) Lebanon, N. H.; Pomfret, Windsor, and Royalton, Vt.; not common.

H. ELLIPTICUM, Hook. Wet places.

H. PERFORATUM, L. (Common St. John's-wort.) Fields; European.

H. MACULATUM, Walt. (*H. CORYMBOSUM*, Muhl.) Damp places; common.

H. MUTILUM, L. Damp places.

H. CANADENSE, L. Very common in sandy soil.

var. *majus*, Gray. Not uncommon.

H. NUDICAULE, Walt. (*H. SAROTHRA*, Michx.)—(Orange Grass.) Sandy fields; common.

Elodes, Adans. Marsh St. John's-wort.

E. CAMPANULATA, Pursh. (*E. VIRGINICA*, Nutt.) Swamps.

ORDER 14. **MALVACEÆ**. MALLOW FAMILY.**Malva**, L. Mallows.

M. ROTUNDIFOLIA, L. (Common Mallow.) Near dwellings.

M. CRISPA, L. (Curled Mallow.) Occasionally seen on the roadsides.

M. MOSCHATA, L. (Musk Mallow.) Roadsides and fields.

M. SYLVESTRIS, L. (High Mallow.) Old gardens; all European.

Abutilon, Tourn. Indian Mallow.

A. AVICENNÆ, Gærtn. (Velvet-Leaf.) Roadsides and waste heaps; from India.

Hibiscus, L. Rose-Mallow.

[*H. TRIONUM*, L. (Flower-of-an-hour.) State Farm, Hanover; occasionally seen; European.]

ORDER 15. **TILIACEÆ**. LINNEN FAMILY.**Tilia**, Tourn. Basswood. Linden. Lime-tree.

T. AMERICANA, L. Woods; common.

[*T. EUROPEA*, L. (European Linden.) Planted by the roadsides and in private grounds. This tree (the LIN) gave the family name to Linnæus.]

ORDER 16. LINACEÆ. FLAX FAMILY.

Linum, Tourn. Flax.

L. Virginianum, L. (Wild Flax.) Dry woods.

L. USITATISSIMUM, L. (Cultivated Flax.) In grain fields. Plymouth, etc.,
Sargent. European.

ORDER 17. GERANIACEÆ. GERANIUM FAMILY.

Geranium, Tourn. Cranesbill.G. maculatum, L. (Cranesbill Geranium.) Hanover, N. H.; Norwich, Vt.;
etc. Open fields.

G. Robertianum, L. (Herb Robert.) Common on moist rocks.

G. Carolinianum, L. Not uncommon; in waste places.

Erodium, L'Her. Storksbill.E. CICUTARIUM, L'Her. Ludlow, Vt., *Miss Pollard*. Rarely met with;
European.**Impatiens**, L. Balsam. Jewel-weed.I. pallida, Nutt. Windsor and Hartland, Vt., etc., *Leland*. Less common
than the next.

I. fulva, Nutt. Common in wet places.

Oxalis, L. Wood-Sorrel.

O. Acetosella, L. (True Wood-Sorrel.) Common.

O. violacea, L. (Violet Wood-Sorrel.) Cavendish, Vt., *Dr. Graves*. Very
unusual so far north, and probably rare.O. corniculata, L. var. stricta, Sav. (O. stricta, L.)—(Yellow Wood-Sor-
rel.) A common weed and very variable.

ORDER 18. RUTACEÆ. RUE FAMILY.

Xanthoxylum, L. Prickly Ash.X. Americanum, Mill. Claremont, N. H., *Dr. Barrows*; Norwich and Hart-
ford, Vt., etc.

[PTELEA TRIFOLIATA, L. (Hop-tree.) From the West, is often planted.]

ORDER 19. ILCINEÆ. HOLLY FAMILY.

(Formerly AQUIFOLIACEÆ.)

Ilex, L. Holly.

I. verticillata, Gray. (Black Alder.) Common in low grounds.

Nemopanthes, Raf. (Mountain Holly.)

N. fascicularis, Raf. (N. Canadensis, DC.) Common in wet cold woods.

ORDER 20. CELASTRACEÆ. STAFF-TREE FAMILY.

Celastrus, L. Staff-tree.

C. scandens, L. (Climbing Bitter-sweet.) Along streams.

[EUONYMUS ATROPURPUREUS, Jacq. (Burning Bush.) From the West;
common in cultivation.]

ORDER 21. RHAMNACEÆ. BUCKTHORN FAMILY.

Rhamnus, Tourn. Buckthorn.

R. alnifolia, L'Her. Bear Pond, Canaan, N. H., J. B. Wallace; Norwich and Sharon, Vt., Jesup; rare.

R. CATHARTICA, L. Used for hedges; sometimes running wild; European.

Ceanothus, L. New Jersey Tea.

C. Americanus, L. Dry woodlands. Lebanon near Conn. river, etc.

ORDER 22. VITACEÆ. VINE FAMILY.

Vitis, Tourn. Grape.

V. Labrusca, L. (Fox grape.) Often near old homesteads, sometimes cultivated; doubtful if native in this region.

V. æstivalis, Michx. (Summer grape.) Possibly native.

V. cordifolia, Michx. As the species is now understood, it may be that plants referred to this species should be included in the next.

V. riparia, Michx. (Frost grape.) Common. This is V. cordifolia, var. riparia, of Gray's Man. (1870). Dr. Engelmann writes (1876):—"I doubt if V. cordifolia grows at all in New England." See his synopsis of the Genus in Am. Nat., Vol. VI, p. 539. See, also, Gray's Manual (1890).

Ampelopsis, Michx. Woodbine. Virginian Creeper.

A. quinquefolia, Michx. Common on river-banks, etc.

ORDER 23. SAPINDACEÆ. SOAPBERRY FAMILY.

Aesculus, L. Horse-chestnut. Buckeye.

[Æ. HIPPOCASTANUM, L. (Horse-chestnut.) Common in cultivation everywhere. Native of Europe.]

[Æ. GLABRA, Willd. (Fetid or Ohio Buckeye.) Not native in New England, but not rare in cultivation along with Æ. PAVIA.]

Acer, Tourn. Maple.

A. Pennsylvanicum, L. (Striped Dogwood. Moose-Wood.) Woods; common.

A. spicatum, Lam. (Mountain Maple.) Common with the above.

A. saccharinum, Wang. (Sugar Maple. Rock Maple.) Very common.

var. nigrum, Torr. & Gray. (Black Sugar Maple.) Probably not uncommon. Recognized by Michaux as a distinct species, and common outside of New England. He noticed it "at Windsor, Vt., but of inferior size." (Michaux's N. A. Sylva, Vol. I, p. 163.) It is seldom distinguished from the ordinary Sugar Maple.

A. dasycarpum, Ehrh. (White or Silver Maple.) On river-banks.

A. rubrum, L. (Red or Swamp Maple.) Very common.

[A. PLATANOIDES (Norway Maple), with fruit having long divergent wings, and A. PSEUDO-PLATANUS with long racemes of pubescent fruit, are often planted.]

Negundo, Moench. Ash-leaved Maple. Box-Elder.

N aceroides, Moench. Not native in New England, but often planted and sometimes spontaneous, as on the White River at Sharon, Vt., Dr. Baxter.

Staphylea, L. Bladder-Nut.

S. trifolia, L. Claremont and Lebanon, N. H.; Norwich, *Sargent*, and Windsor, Vt. River-banks; not very common.

ORDER 24. **ANACARDIACEÆ**. CASHEW FAMILY.**Rhus**, L. Sumach.

R. typhina, L. (Staghorn Sumach.) Fields; very common. This species and the next two are not poisonous.

R. glabra, L. (Smooth Sumach.) Fields; common.

R. copallina, L. Claremont, N. H., *Dr. Barrows*; Holderness, N. H., and Fairlee, Vt., *Jesup*. Rare.

R. venenata, DC. (Poison Sumach. Poison Dogwood.) Claremont, N. H., *Dr. Barrows*; Mud Pond, Canaan, N. H., *Jesup*; Windsor, Vt., *Leland*. Not uncommon in swamps throughout the district.

R. Toxicodendron, L. (Poison Ivy.) Very common, and, like the preceding, very poisonous to the touch.

ORDER 25. **POLYGALACEÆ**. MILKWORT FAMILY.**Polygala**, Tourn. Milkwort.

P. paucifolia, Willd. (Fringed Polygala.) Borders of woods; common.

P. polygama, Walt. Claremont, *Dr. Barrows*; New London, *Miss Messer*.

P. sanguinea, L. Charlestown and New London, N. H.; Windsor, Vt.; etc.

P. verticillata, L. Claremont, *Dr. Barrows*.

var. *ambigua* (*P. ambigua*, Nutt.) Lebanon, *Sargent*.

ORDER 26. **LEGUMINOSÆ**. PULSE FAMILY.**Lupinus**, Tourn. Lupine.

L. perennis, L. Claremont, etc., *Dr. Barrows*; Plymouth, *Miss Langdon*; rare; abundant farther south.

Trifolium, Tourn. Clover. Trefoil.

T. ARVENSE, L. (Stone Clover.) Old fields and roadsides.

T. PRATENSE, L. (Red Clover.) Meadows.

T. repens, L. (White Clover.) Everywhere common.

T. AGRARIUM, L. (Yellow or Hop-Clover.) Meadows and roadsides.

T. PROCUMBENS, L. Claremont, *Dr. Barrows*; Royalton, Vt., *H. Q. Ward*. Uncommon; has pinnately trifoliate leaves and very small heads. This and all the preceding species are European, except *T. repens*.

T. HYBRIDUM, L. (Alsike or Dutch Clover.) From Holland. Meadows.

Melilotus, Tourn. Melilot.

M. OFFICINALIS, Willd. (Yellow Melilot.) Waste or cultivated grounds; both species from Europe.

M. ALBA, Lam. (White Melilot or Sweet Clover.) Near dwellings and sometimes in gardens.

Medicago, Tourn. Medick.

- M. SATIVA, L. (Lucerne, Alfalfa.) Claremont, N. H., *Dr. Barrows*; Pomfret, Vt., *A. P. Morgan*. Cultivated sometimes for green fodder.
- M. LUPULINA, L. (Black Medick.) Claremont, *Dr. Barrows*. In grass ground, Hanover, etc., and often mistaken for a *Trifolium*.
- M. MACULATA, Willd. (Spotted Medick.) Claremont, *Dr. Barrows*. This and the next species introduced in foreign wool, and probably transient visitors. The entire genus is European.
- M. DENTICULATA, Willd. Claremont, *Dr. Barrows*.

Robinia, L. Locust-tree.

- R. Pseudacacia, L. (Common Locust.) Naturalized from Penn. and the Southwest.
- R. viscosa, Vent. (Clammy Locust.) Native of Virginia, and spreading freely.

Astragalus, Tourn. Milk-Vetch.

- A. alpinus, L. On rocks at Sumner's Falls, Plainfield, N. H., *Flint*: very rare.
- A. Robbinsii, Gray. Sumner's Falls, N. H., *Flint*; Hartland, Vt., *Leland*; very rare.

Desmodium, Desv. Tick-Trefoil.

- D. nudiflorum, DC. Dry woods; common.
- D. acuminatum, DC. Woods; common.
- D. rotundifolium, DC. Dry rocky woods.
- D. Dillenii, Darl. Open woodlands.
- D. paniculatum, DC. Woods.
- D. Canadense, DC. Dry woods; common.
- D. rigidum, DC. Dry hillsides.
- D. Marilandicum, Boott. Copses; not very common.

Lespedeza, Michx. Bush-Clover.

- L. violacea, Pers. Dry woods.
- L. polystachya, Michx. (L. hirta, Ell.) Dry hillsides; common.
- L. capitata, Michx. Dry hillsides; frequent.

Vicia, Tourn. Vetch or Tare.

- V. SATIVA, L. (Common Vetch.) Cultivated fields; not uncommon; from Europe.
- VAR. ANGUSTIFOLIA, Seringe. Hanover.
- V. Cracca, L. Infrequent; usually in grass land.

Apios, Boerhaave. Ground-nut.

- A. tuberosa, Moench. Windsor, Vt., *Flint*; Royalton, Vt., *L. Wild*; Plymouth, N. H., *Sargent*.

Amphicarpæa, Ell. Hog Pea-nut.

- A. monoica, Nutt. Borders of woods; very common.

Cassia, Tourn. Wild Senna.

- C. Marilandica, L. Hartford, Vt. Alluvial soil; rare.

Gleditschia, L. Honey-Locust.

G. triacanthos, L. Never truly wild in New England; common in cultivation; from the Western States, and sometimes thornless.

[Additional plants of this order:—*CORONILLA VARIA* (Purple Coronilla), *ARACHIS HYPOGAEA* (Pea-nut), *CARAGANA ARBORESCENS* (Pea-tree), *WISTARIA SINENSIS* (Chinese Wistaria), *PHASEOLUS* (Garden Bean), *PISUM SATIVUM* (Garden Pea), besides others from the Western States and from Europe.]

ORDER 27. **ROSACEÆ. ROSE FAMILY.****Prunus**, Tourn. Plum. Cherry.

P. Americana, Marshall. (Wild Yellow or Red Plum.) Not common.

P. pumila, L. (Dwarf Cherry.) Islands in the Conn. river near Hanover, etc. *Jesup.*

P. Pennsylvanica, L. (Red Cherry.) Very common everywhere.

P. Virginiana, L. (Choke-Cherry.) Very common.

P. serotina, Ehrh. (Black Cherry.) Common.

Spiræa, L. Meadow-Sweet.

S. salicifolia, L. (Common Meadow-Sweet.) Roadsides and pastures.

S. tomentosa, L. (Hardhack. Steeple-Bush.) Roadsides; common.

S. SORBIFOLIA, from Siberia, with leaves like the Mountain Ash, is well established in many places.

Rubus, Tourn. Raspberry. Blackberry.

R. odoratus, L. (Purple Flowering Raspberry.) Often wrongly called Mulberry.

R. triflorus, Richardson. (Dwarf Raspberry.) Much like a Blackberry.

R. strigosus, Michx. (Wild Red Raspberry.)

R. occidentalis, L. (Black Raspberry. Thimbleberry.) Both this and all the preceding species common.

R. villosus, Ait. (High Blackberry.) Very variable in size and shape of fruit. Very common.

R. Canadensis, L. (Low Blackberry.) Dry fields. The smaller and trailing forms of the last connect with this species.

R. hispida, L. (Running Swamp-Blackberry.) Not very common.

Dalibarda, L.

D. repens, L. Low woods; common.

Geum, L. Avens.

G. album, Gmelin. Borders of woods; common.

G. Virginianum, L. Borders of woods; common.

G. macrophyllum, Willd. Killington Mt., Vt., *Rev. H. M. Denslow.*

G. strictum, Ait. Moist meadows; frequent.

G. rivale, L. (Water Avens.) Wet meadows.

Waldsteinia, Willd.

W. fragarioides, Tratt. (Barren Strawberry.) Wooded hillsides; common.

Fragaria, Tourn. Strawberry.

F. Virginiana, Mill. (Field Strawberry.) Everywhere common.

F. vesca, L. (Wood Strawberry.) Less common than the preceding.

Potentilla, L. Cinque-foil. Five-Finger.

P. arguta, Pursh. Roadsides. Hanover; Lebanon; not rare.

P. Norvegica, L. Dry fields; common.

P. argentea, L. (Silvery Cinque-foil.) Dry, barren fields; common.

P. palustris, Scop. (Marsh Five-Finger.) Cold bogs; George Pond, Enfield, etc.

P. fruticosa, L. (Shrubby Cinque-foil.) Claremont, N. H., *Dr. Barrows*; Chittenden, Vt., *Flint*; a shrub.

P. tridentata, Ait. (Three-toothed Cinque-foil.) Kearsarge Mt., *Dr. Barrows*; Moosilauke Mt., *Flint*.

P. Anserina, L. (Silver-weed.) Claremont, N. H., *Dr. Barrows*; Royalton, Vt., *Ward*. Near ponds and streams.

P. Canadensis, L. (Common Five-Finger.) Dry soil; very common; stem commonly prostrate; sometimes erect.

Agrimonia, Tourn. Agrimony.

A. Eupatoria, L. (Common Agrimony.) Roadsides.

Poterium, L. Burnet.

P. Canadense, Benth & Hook. (Canadian Burnet.) Bristol, N. H., etc., *F. L. Pattee*. Not common.

Rosa, Tourn. Rose.

R. blanda, Ait. (Early Rose.) Banks of Conn. river, Hanover, N. H.; Fairlee, Vt., *Sargent*. Not very common.

R. Carolina, L. (Swamp Rose.) Swamps and river-banks throughout.

R. humilis, Marsh. (*R. lucida* of most authors.)—(Dwarf Rose.) With the same range as *R. blanda*, but more common. The true *lucida* is found on the New Hampshire seacoast.

R. nitida, Willd. (Shining Rose.) New London, *Sargent*; George Pond, Enfield, etc., *Jesup*. Rather rare.

R. RUBIGINOSA, L. (Sweet-brier.) Roadsides and pastures; not abundant, but widely distributed; European.

R. CINNAMOMEA, L. (Cinnamon Rose.) Native of Europe, very common about old dwellings, always double.

Pyrus, L. Pear. Apple.

P. arbutifolia, L. (Choke-berry.) Swamps; common, with red fruit.

var. *melanocarpa*, Hook. With black fruit.

P. Americana, DC. (Mountain Ash.) Common.

Crataegus, L. White Thorn. Hawthorn.

[*C. OXYACANTHA*, L. (English Hawthorn.) Cultivated for ornamental purposes, and sometimes spontaneous.]

C. coccinea, L. (Scarlet-fruited Thorn.) Common in hillside pastures, etc. var. *macracantha*, Dudley. Frequent and quite distinct from the type.

Perhaps a distinct species. Hanover, Lyme, Lebanon, and Enfield, N. H.; Norwich, Hartford, etc., Vt.

C. punctata, Jacq. Widely distributed on both sides of the Conn. river.

C. Crus-galli, L. Claremont, *Dr. Barrows*; College Park, Hanover. Not common, and possibly not indigenous.

Amelanchier, Medic. June-berry.

A. Canadensis, Torr. & Gray. (var. *Botryapium*, T. & G.)—(Shad-bush.)

The more common form.

var. *oblongifolia*, Gray. Not rare; two weeks later. A smaller tree or shrub.

[This Order also includes numerous species of the Genus *PRUNUS* (Plum, Cherry, Peach, Nectarine, and Apricot); *SPIRÆA* (many species); *PYRUS COMMUNIS* (Pear); *PYRUS MALUS* (Apple); *PYRUS PRUNIFOLIA* (Siberian Crab-apple); and *CYDONIA VULGARIS* (Common Quince).]

ORDER 28. **SAXIFRAGACEÆ.** SAXIFRAGE FAMILY.**Saxifraga**, L. Saxifrage.

S. Virginiana, Michx. (Rock Saxifrage.) Common on rocks.

S. Pennsylvanica, L. (Swamp Saxifrage.) Common in swamps.

Tiarella, L. False Mitre-wort.

T. cordifolia, L. Common in moist ground.

Mitella, Tourn. Mitre-wort. Bishop's-Cap.

M. diphylla, L. Hillsides in rich woods; very common.

M. nuda, L. Wet woods in moss. Hanover, N. H.; Norwich, Vt.; etc.
Not rare.

Chrysosplenium, Tourn. Golden Saxifrage.

C. Americanum, Schwein. Common in wet places.

Parnassia, Tourn. Grass of Parnassus.

P. Caroliniana, Michx. Queechy Gulf in Hartland, Norwich, and Royalton, Vt., *L. Wild.* Not common.

Philadelphus, L. Mock Orange. Syringa.

[*P. GRANDIFLORUS*, Willd. Cultivated from Virginia.]

[*P. CORONARIUS*, L. In cultivation from Europe.]

Ribes, L. Currant. Gooseberry.

R. Cynosbati, L. (Prickly Gooseberry.) Common.

R. oxyacanthoides, L. (*R. hirtellum*, Michx.)—(Smooth Gooseberry.) Meriden and Hanover, N. H.; Union Village and Sharon, Vt. Probably not rare in swamps.

R. lacustre, Poir. Hanover and Meriden, N. H.; Mt. Ascutney, Vt. Cold woods and swamps.

R. prostratum, L'Her. (Fetid Currant.) Common in damp, rocky woods.

R. floridum, L'Her. (Wild Black Currant.) Woods, and frequent in old gardens.

R. rubrum, L. var. *subglandulosum*, Maxim. This is the native Red Currant of the White Mts. and Northern New England. The Red Currant of the garden is the European form (*R. rubrum*) introduced by the early settlers. The two forms however cannot readily be distinguished.

R. aureum, Pursh. (Missouri Currant.) Common in cultivation, and persisting for years about old homesteads. Native of Missouri and the West.

ORDER 29. CRASSULACEÆ. ORPINE FAMILY.

Penthorum, Gronov. Ditch Stone-crop.

P. sedoides, L. Open wet places.

Sedum, Tourn. Stone-crop.

S. ACRE, L. (Mossy Stone-crop.) Norwich, Chelsea, and Royalton, Vt.; on rocks, etc. Frequent in gardens throughout the district.

S. TELEPHIUM, L. (Live-for-ever.) Very widely distributed and thoroughly naturalized; both species European.

ORDER 30. DROSERACEÆ. SUNDEW FAMILY.

Drosera, L. Sundew.

D. rotundifolia, L. Peat bogs and banks of brooks; common.

D. intermedia, Hayne. var. Americana, DC. (D. longifolia. Man., 5th ed.) Hanover, Canaan, Enfield, Plainfield, etc. Sandy shores of ponds and on decaying logs.

ORDER 31. HAMAMELIDÆ. WITCH-HAZEL FAMILY.

Hamamelis, L. Witch-Hazel.

H. Virginiana, L. Peculiar for blooming in October and not perfecting its fruit until the next autumn; common in woods.

ORDER 32. HALORAGEÆ. WATER-MILFOIL FAMILY.

Myriophyllum, Vaill. Water-Milfoil.M. spicatum, L. Enfield Pond, *Jesup*; not rare.M. verticillatum, L. Mascoma Creek, West Lebanon, etc., *Jesup*.M. tenellum, Bigel. Shores of Enfield Pond, *Jesup*; not rare.**Proserpinaca**, L. Mermaid-weed.

P. palustris, L. Swamps and ditches; common.

Hippuris, L. Mare's Tail.

[H. vulgaris, L. About ponds; rare. Near Long Pond, West Danville, Vt., Dr. Blanchard, and in Mass.; probably within our limits.]

Callitrichæ, L. Water-Starwort.

C. verna, L. Ponds; common.

ORDER 33. LYTHRACEÆ. LOOSESTRIFE FAMILY.

Lythrum, L. Loosestrife.

L. SALICARIA, L. (Spiked Loosestrife.) Claremont, N. H., Dr. Graves; Sharon, Vt., etc.; often cultivated, and commonly trimorphous.

ORDER 34. ONAGRACEÆ. EVENING-PRIMROSE FAMILY.

Ludwigia, Ell. False Loosestrife.

L. palustris, L. (Water Purslane.) Ditches; common.

Epilobium, L. Willow-herb.

E. angustifolium, L. (Often like Erechtites, called "Fire-weed.") Common in moist and newly cleared lands.

E. lineare, Muhl. (*E. palustre*, var. *lineare*, Man., 5th ed.) Bogs; common.
E. strictum, Muhl. (*E. molle*, Torr.) Claremont, *Dr. Barrows*; not common.

E. coloratum, Muhl. Wet places; very common.

[A number of new species of this genus have recently (Man., 6th ed.) been assigned to this region, some of which doubtless are collected as forms of the above and remain to be distinguished from them.]

Oenothera, L. Evening Primrose.

O. biennis, L. Fields and roadsides everywhere; many vars.; one with large flowers is found in gardens (var. *grandiflora*).

O. pumila, L. Dry fields; common.

Circæa, Tourn. Enchanter's Nightshade.

C. Lutetiana, L. Damp shades; common.

C. alpina, L. Rocky, wet woodlands; much like a small form of the former.

ORDER 35. **CUCURBITACEÆ**. CUCUMBER FAMILY.

Sicyos, L. Bur-Cucumber.

S. angulatus, L. Wet, shady places and waste heaps; a not uncommon weed in Hanover, etc.

Echinocystis, Torr. & Gray. Balsam-Apple.

E. lobata, Torr. & Gray. Banks of streams, Hanover, N. H.; Sharon, Vt.; etc.

[Cultivated species of this Order belong to the Genera:—*LAGENARIA* (Bottle Gourd), *CUCURBITA* (Pumpkin, Squash, and Egg Gourd), *CITRULLUS* (Watermelon), *CUCUMIS* (Melon and Cucumber).]

ORDER 36. **FICOIDEÆ**.

Mollugo, L. Carpet-weed.

M. verticillata, L. A common weed in cultivated ground. Formerly included in the Pink Family (CARYOPHYLLACEÆ).

ORDER 37. **UMBELLIFERÆ**. PARSLEY FAMILY.

Daucus, Tourn. Wild Carrot.

D. CAROTA, L. A noxious weed of the old world; the cultivated carrot run wild.

Angelica, L.

A. atropurpurea, L. (*Archangelica atropurpurea*, Hoffm.) River-banks.

Conioselinum, Fisch. Hemlock-Parsley.

C. Canadense, Torr. & Gray. Near Goose Pond, Canaan.

Heracleum, L. Cow-Parsnip.

H. lanatum, Michx. Moist, rich ground; common.

Pastinaca, L. Wild Parsnip.

P. SATIVA, L. Fields; common; from Europe.

Ligusticum, L. Lovage.

L. Scoticum, L. Near dwellings; Hanover, etc.; perhaps a garden escape.

Cryptotaenia, DC. Honewort.

C. Canadensis, DC. Thickets; not rare.

Sium, Tourn. Water-Parsnip.

S. cicutaeifolium, Gmelin. (S. lineare, Michx.) Wet places; common.

Zizia, Koch. Meadow-Parsnip.

Z. aurea, Koch. (Thaspium aureum, var. apterum, Gray's Man., 5th. ed.) In moist meadows; common.

Carum, L. Caraway.

C. CARUM, L. Everywhere near dwellings; European.

Cicuta, L. Water-Hemlock.

C. maculata, L. (Spotted Cowbane.) Root very poisonous; common.

C. bulbifera, L. Swamps; seldom fruiting, the flowers replaced by bulb-lets.

Conium, L. Poison Hemlock.

C. MACULATUM, L. On waste heaps; a virulent poison. Hanover, etc. From Europe.

Osmorrhiza, Raf. Sweet Cicely.

O. brevistylis, DC. Rich, moist woods.

O. longistylis, DC. With, but less common than, the preceding.

Hydrocotyle, Tourn. Water Pennywort.

H. Americana, L. Wet places; common.

Sanicula, Tourn. Sanicle. Black Snakeroot.

S. Marilandica, L. With the variety but more common. Borders of woods.

var. Canadensis, Torr. (S. Canadensis, L.)

[This large Order also includes,—APIUM GRAVEOLENS (Celery), CARUM PETROSELINUM (Parsley), FENICULUM VULGARE (Fennel), etc.]

ORDER 38. **ARALIACEÆ. GINSENG FAMILY.****Aralia**, Tourn.

A. racemosa, L. (Spikenard.) Rich woodlands; often cultivated.

A. hispida, Vent. (Bristly Sarsaparilla.) Rocky places; frequent.

A. nudicaulis, L. (Wild Sarsaparilla.) Borders of woods; common.

A. quinquefolia, Dec. & Pla. (Ginseng.) Rich woods; not infrequent.

A. trifolia, Dec. & Pla. (Dwarf Ginseng. Ground-nut.) Very common in woods.

ORDER 39. **CORNACEÆ. DOGWOOD FAMILY.****Cornus**, Tourn. Dogwood. Cornel.

C. Canadensis, L. (Bunch-berry.) Common.

C. florida, L. (Flowering Dogwood.) Plainfield, N. H., Miss Stevens; Springfield, Vt., Rev. S. Hayward; Charlestown and New London, N. H.; Pomfret, Vt., etc.; rare so far northward.

- C. circinata, L'Her. (Round-leaved Cornel.) Hanover, N. H.; West Hartford, Vt., *Flint*; and Killington Mt., Vt., *Rev. H. M. Denslow*.
C. sericea, L. (Silky Cornel.) Borders of ponds; frequent.
C. stolonifera, Michx. (Red-osier Dogwood.) Wet places; common.
C. paniculata, L'Her. (Panicled Cornel.) Charlestown, *Dr. Graves*; Plymouth, *Miss Langdon*: rare.
C. alternifolia, L. (Alternate-leaved Cornel.) Common by the roadside everywhere.

Nyssa, L. Tupelo. Pepperidge. Sour-Gum Tree.

N. sylvatica, Marsh. (N. multiflora, Wang.) On the outlet of Squam Lake, N. H.; not common. [Not rare in Boscowen, *Barton*.]

DIVISION II. GAMOPETALOUS EXOGENS.

ORDER 40. CAPRIFOLIACEÆ. HONEYSUCKLE FAMILY.

Sambucus, Tourn. Elder.*S. Canadensis*, L. (Black-berried Elder.) Common.*S. racemosa*, L. (*S. pubens*, Michx.)—(Red-berried Elder.) Common.**Viburnum**, L. Arrow-wood.*V. lantanaoides*, Michx. (Hobble-bush.) Frequent in woods.*V. Opulus*, L. (High Cranberry-bush.) Hanover, Lebanon, and New London, N. H.; Sharon, etc., Vt.; low ground, not rare. The Snow Ball Tree from Europe is a cultivated form of this species.*V. pauciflorum*, Pylaie. Claremont, *Dr. Barrows*; belongs farther northward, and doubtless rare.*V. acerifolium*, L. (Maple-leaved Arrow-wood.) Rocky woods; common.*V. dentatum*, L. (Arrow-wood.) Wet places.*V. cassinoides*, L. (*V. nudum*, var. *cassinoides*, Torr. & Gray.)—(Witherod.) Swamps in Lebanon and Hanover, N. H.; Sharon and Hartford, Vt.; etc.; common.*V. Lentago*, L. (Sheep-berry.) Wet river-banks, etc.; not rare.**Triosteum**, L. Fever-wort. Horse-Gentian.*T. perfoliatum*, L. White River Junction and Norwich, Vt., etc. Rich woodlands.**Linnæa**, Gronov. Twin-flower.*L. borealis*, Gronov. Common in bogs and mossy woods. The favorite flower of the great Linnæus, and named for him.**Symporicarpos**, Dill. Snowberry.[*S. racemosus*, Michx. Common in cultivation; native of western Vt. *S. VULGARIS* and *S. OCCIDENTALIS* from the Western States are sometimes in private grounds.]**Lonicera**, L. Honeysuckle. Woodbine.*L. ciliata*, Muhl. (Fly-Honeysuckle.) Rocky woods; common.*L. cœrulea*, L. (Mountain Fly-Honeysuckle.) Claremont, *Dr. Barrows*; Bear Pond, Canaan, *E. S. Comings*; Enfield Pond, *A. C. French*.*L. glauca*, Hill. (*L. parviflora* of Man., 5th ed.) (Twining Honeysuckle.) Hanover, N. H.; Norwich, Vt.; etc.; not rare.[*L. TATARICA*, L. (Tartarian Honeysuckle.) The common Bush Honeysuckle of cultivation; from Europe.]**Diervilla**, Tourn.*D. trifida*, Moench. (Bush Honeysuckle.) Common in fence rows.

ORDER 41. RUBIACEÆ. MADDER FAMILY.

Houstonia, L.*H. cœrulea*, L. (Bluets. Innocence.) Pastures; commonly dimorphous.

Cephalanthus, L. Button-bush.

C. occidentalis, L. New Found Lake, *Dr. Barrows*; Goose Pond, Canaan; Andover, *A. C. French*; Plymouth, *Sargent*; etc.

Mitchella, L. Partridge-berry.

M. repens, L. Dry woods; common under trees and dimorphous.

Galium, L. Bedstraw. Cleavers.

G. MOLLUGO, L. North Pomfret, Vt., *Morgan*.

G. Aparine, L. (Cleavers. Goose-Grass.) Moist grounds, Norwich, Vt., *Eggleson*; not common.

G. Kamtschaticum, Steller, (*G. circæzans*, var. *montanum*, Torr. & Gray.) Killington Peak, Vt., *Denslow*.

G. circæzans, Michx. (Wild Liquorice.) Woods; not rare.

G. lanceolatum, Torr. (Wild Liquorice.) Pomfret, Vt., *Morgan*; Hanover, *French*,—also Weirs, N. H.; infrequent in dry woods.

G. trifidum, L. (Small Bedstraw.) Swamps; very variable; common.

G. asprellum, Michx. (Rough Bedstraw.) Low thickets; abundant.

G. triflorum, Michx. (Sweet-scented Bedstraw.) Rich woodlands; common.

ORDER 42. **COMPOSITÆ. COMPOSITE FAMILY.****Eupatorium**, Tourn. Thoroughwort. Boneset.

E. purpureum, L. (Purple Thoroughwort.) Common in swamps.

E. perfoliatum, L. (Medicinal Thoroughwort.) Low grounds; common.

E. ageratoides, L. (White Snake-root.) Rich woods.

Solidago, L. Golden-rod.

S. squarrosa, Muhl. Claremont, N. H., *Graves*: Windsor, *Flint*; Killington Mt., *Eggleson*, and Fairlee Pond, Vt., *Jesup*.

S. cæsia, L. Borders of woodlands; common.

S. latifolia, L. Shaded banks; common.

S. bicolor, L. Dry banks. Also var. *concolor*, Torr. & Gray, yellow rays.

S. macrophylla, Pursh. (*S. thyrsoidea*, E. Meyer.) Claremont, N. H., *Dr. Barrows*; Mt. Ascutney, *Leland*,—and Killington Mt., Vt., *Eggleson*.

S. Virgaurea, L., var. *alpina*, Bigel. Moosilauke Mt., *Flint*; Killington Peak, Vt., *Dr. Robbins*.

S. humilis, Pursh. (*S. Virgaurea*, var. *humilis*, Gray.) Mountains.

S. puberula, Nutt. Killington, Mt., Vt., *Denslow*; Rochester, Vt., *Eggleson*.

S. uliginosa, Nutt. (*S. stricta*, Ait.) Peat bogs. Canaan and Hanover, N. H.; Sharon, *Dr. Baxter*, and Norwich, Vt., *Jesup*: occasional.

S. rugosa, Mill. (*S. altissima*, Man., 5th ed.) Fields; common.

S. neglecta, Torr. & Gray. Swamps, and very variable. Some forms as determined by Dr. Gray are much like "uliginosa" and "arguta." Perhaps not very rare: Canaan, *Jesup*.

S. arguta, Ait. (*S. Muhlenbergii*, Torr. & Gray.) Borders of woods; frequent.

S. juncea, Ait. (*S. arguta*, Torr. & Gray.) Borders of woodlands; common.

- S. serotina, Ait. (*S. gigantea*, Man., 5th ed.) Roadsides; common.
 var. *gigantea*, Gray. (*S. serotina*, Man., 5th ed.) low ground; common.
S. Canadensis, L. Fields and roadsides; very common and variable.
S. nemoralis, Ait. Dry, sterile fields; very common.
S. lanceolata, L. Roadsides, etc.; very common.

Sericocarpus, Nees. White-topped Aster.

- S. conyzoides*, Nees. Dry grounds.

Aster, L. Starwort. Aster.

- A. corymbosus*, Ait. Woodlands; common.
A. macrophyllus, L. Borders of woods; common.
A. Novæ-Angliæ, L. Royalton, Vt., *Ward*; more common southward.
A. patens, Ait. Dry ground.
A. undulatus, L. Dry woods; common.
A. cordifolius, L. Woodlands; very common.
 [A. *Lindleyanus*, T. & G. has been found at Lisbon, N. H., by C. E. *Faxon*, just beyond our limits.]
A. laevis, L. Woodlands.
A. ericoides, L. *Dr. Barrows*. More common southward.
A. polyphyllus, Willd. "Heads larger and flowering earlier than the preceding."
A. multiflorus, Ait. Claremont, N. H., *Dr. Graves*: Norwich, Vt., *Sargent*.
A. dumosus, L. Claremont, *Dr. Barrows*; scarce.
A. vimineus, Lam. (*A. Tradescanti*, Man., 5th ed.) Moist banks and variable.
A. diffusus, Ait. (*A. miser*, Man., 5th ed.) Roadsides and fields; very common and variable.
A. Tradescanti, L. (*A. tenuifolius*, Man., 5th ed.) With the next, but not readily distinguished from it.
A. paniculatus, Lam. (*A. simplex*, Man., 5th ed.) Moist banks; very common.
A. salicifolius, Ait. (*A. carneus*, Man., 5th ed.) What appears to be this species has been found with the preceding.
A. Novi-Belgii, L. (*A. longifolius*, Man., 5th ed.) Moist ground and very variable.
A. puniceus, L. Swamps; very common and variable.
A. umbellatus, Mill. (*Diplopappus umbellatus*, Torr. & Gray.) Moist thickets; common.
A. linariifolius, L. (*Diplopappus linariifolius*, Hook.) Dry grounds; not very common.
A. acuminatus, Michx. Rich woods; frequent.
A. nemoralis, Ait. George Pond, Enfield. [Todd Pond, Bradford, *Barton*.]

Erigeron, L. Fleabane.

- E. Canadensis*, L. (Horseweed.)
E. annuus, Pers. (Daisy Fleabane.) Fields and waste places.
E. strigosus, Muhl. (Daisy Fleabane.) A coarser plant than the rest.
 This and both the above common weeds.

- E. hyssopifolius, Michx. (*Aster graminifolius*, Pursh.) On rocks in Quechee Gulf, Vt., *Jesup*: quite rare.
 E. bellidifolius, Muhl. (Daisy-leaved Fleabane. Robin's Plantain.) Moist meadows and banks; very common.
 E. Philadelphicus, L. Moist ground; common.

Antennaria, Gaertn. Plantain-leaved Everlasting.

- A. plantaginifolia, Hook. Sterile banks; very common.

Anaphalis, DC. Pearly Everlasting.

- A. margaritacea, Benth. & Hook. Dry fields; very common.

Gnaphalium, L. Cudweed.

- G. polycephalum, Michx. (Common Everlasting.)
 G. decurrens, Ives. (Everlasting.) Old fields.
 G. uliginosum, L. (Low Cudweed.) Roadsides. All the species common.

Inula, L. Elecampane.

- I. HELENIUM, L. A coarse plant on roadsides and in pastures. Native of Europe.

Ambrosia, Tourn. Ragweed.

- A. trifida, L. (Great Ragweed.) Grantham, *Barton*.
 A. artemisiæfolia, L. (Roman Wormwood.) A common farm weed.

Xanthium, Tourn. Cocklebur. Clotbur.

- X. SPINOSUM, L. On waste heaps. From Tropical America.
 X. STRUMARIUM, L. White River Junction, Vt., *Flint*. Barnyards, etc.; not common. From India.

Heliopsis, Pers. Ox-eye.

- H. scabra, Dunal. (H. levis, var. scabra, T. & G.) Norwich, Vt., *Sargent*. A chance immigrant from the west.

Rudbeckia, L. Cone-flower.

- R. laciniata, L. Low grounds.
 R. hirta, L. Introduced into meadows from the Western States.

Helianthus, L. Sunflower.

- H. giganteus, L. Randolph, Vt., *Bates*; Royalton, Vt., *Ward*.
 H. annuus, L. (Common Sunflower.) Long cultivated and often spontaneous.
 H. strumosus, L. Charlestown, *Dr. Graves*; common farther south.
 H. divaricatus, L. Not common.
 H. decapetalus, L. Banks of streams, etc.; very common.
 H. tuberosus, L. (H. doronicoides, Man., 5th ed.)—(Jerusalem Artichoke.) Often found in old gardens. Dr. Gray has shown that both this species and H. annuus originated in America.

Bidens, L. Bur-Marigold.

- B. frondosa, L. (Common Beggar-ticks.) The entire genus is found in swamps and wet places.
 B. connata, Muhl. (Swamp Beggar-ticks.) Common in ditches, etc.
 B. cernua, L. (Smaller Bur-Marigold.) Not well distinguished from the next; both probably common.

B. chrysanthemoides, Michx. (Larger Bur-Marigold.)

B. Beckii, Torr. (Water Marigold.) Claremont, N. H., Dr. Barrows; Fairlee Pond, Vt., Jesup. Rare.

Galinsoga, Ruiz & Pavon.

G. PARVIFLORA, Cav. From South America. On waste heaps; Windsor, Vt., Leland.

Anthemis, L. Chamomile.

A. COTULA, DC. (Maruta Cotula, DC.)—(May-weed.) From Europe; common by roadsides.

A. ARVENTIS, L. (Corn Chamomile.) From Europe. Occasionally introduced with grass seed.

[A. NOBILIS, L. (Garden Chamomile.) In old gardens.]

Achillea, L. Yarrow.

A. Millefolium, L. Fields; common.

[A. PTARMICA, L. (Sneezewort.) In old gardens; both single and double.]

Chrysanthemum, Tourn. Ox-eye Daisy.

C. LEUCANTHEMUM, L. (Leucanthemum vulgare, Lam.)—(Ox-eye or White Daisy. White-weed.) From Europe. Everywhere in grass ground.

[C. PARTHENIUM, Pers. (Feverfew.) In old gardens.]

Tanacetum, L. Tansy.

T. VULGARE, L. Roadsides everywhere; European weed.

Artemesia, L. Wormwood.

A. VULGARIS, L. (Mugwort.) Near dwellings; from Europe.

A. ABSINTHIUM, L. (Wormwood.) Escaped from cultivation; Enfield, etc.; from Europe.

Tussilago, Tourn. Coltsfoot.

T. FARFARA, L. Clayey soil, along brooks. Thought to have come from Europe, but now found far from cultivation.

Petasites, Tourn. Sweet Coltsfoot.

[P. palmata, Gray. (Nardosmia palmata, Hook.) A rare plant reported on both our northern and southern borders, and probably within our limits.]

Senecio, Tourn. Groundsel.

S. VULGARIS, L. In cultivated grounds; not common; from Europe.

S. aureus, L. (Golden Ragwort.) Several varieties; in both wet and dry ground.

Erechtites, Raf. Fireweed.

E. hieracifolia, Raf. Clearings that have been burned over. This and Epilobium augustifolium are the most common Fireweeds.

Arctium, L. Burdock.

A. LAPPA, L. (Lappa officinalis, All.) Roadsides; very common and variable; from Europe.

Cnicus, Tourn. Thistle.

- C. LANCEOLATUS, Hoffm. (*Cirsium*, Scop.)—(Common Thistle.) From Europe. In pastures.
 C. altissimus, Willd., var. *discolor*, Gray. (*Cirsium discolor*, Spreng.) Fields, etc.
 C. muticus, Pursh. (*Cirsium*, Michx.)—(Swamp Thistle.) Common.
 C. pumilus, Torr. (*Cirsium*, Spreng.)—(Pasture Thistle.) Fields; common.
 C. ARVENSIS, Hoffm. (*Cirsium*, Scop.)—(Canada Thistle.) Everywhere. Introduced from Europe, by way of Canada, during the Revolutionary war.

Krigia, Schreber. Dwarf Dandelion.

- K. Virginica, Willd. Hanover, N. H.; Olcott, Vt., etc.

Cichorium, Tourn. Chicory or Succory.

- C. INTYBUS, L. Fields and roadsides. Hanover, etc. From Europe.

Tragopogon, L. Goat's-beard.

- T. PORRIFOLIUS, L. (Salsify. Oyster-plant.) New London, *Sargent*; Hanover, etc. A garden escape.

Hieracium, Tourn. Hawkweed.

- H. AURANTIACUM, L. From Europe. Found several times by the roadside; Norwich, Vt.; Royalton, Vt., *Ward*.
 H. Canadense, Michx. Dry woods; frequent.
 H. paniculatum, L. Open woods; not uncommon.
 H. venosum, L. (Rattlesnake-weed.) Woods and plains; common.
 H. scabrum, Michx. Dry woods; very common.
 H. Gronovii, L. Not common. *Dr. Barrows*.

Prenanthes, Vaill. Rattlesnake-root.

- P. alba, L. (*Nabalus*, Hook.)—(White Lettuce.) Borders of rich woods.
 P. serpentaria, Pursh. (*Nabalus Fraseri*, DC.)—(Lion's-foot.) Borders of woods; not rare.
 P. altissima, L. (*Nabalus*, Hook.) Rich, moist woods.

Taraxacum, Haller. Dandelion.

- T. OFFICINALE, Weber. (T. *Dens-leonis*, Desf.) Fields and pastures; introduced from Europe.

Lactuca, Tourn. Wild Lettuce.

- L. Canadensis, L. Borders of fields; common.
 L. integrifolia, Bigel. (L. *Canadensis*, var. *integrifolia*, Torr. & Gray.) With the above.
 L. hirsuta, Muhl. (L. *Canadensis*, var. *sanguinea*, Torr. & Gray.) Not uncommon.
 L. leucophæa, Gray. (*Mulgedium*, DC.)—(Blue Lettuce.) Low grounds; common.

Sonchus, L. Sow-Thistle.

- S. OLERACEUS, L. Waste heaps; common.
 S. ASPER, Vill. North Pomfret, Vt., *A. P. Morgan*. Both European weeds.

ORDER 43. LOBELIACEÆ. LOBELIA FAMILY.

Lobelia, L.

- L. cardinalis*, L. (Cardinal-flower.) Borders of streams and ponds in Lebanon, Canaan, New London, Hanover, etc., N. H.; and also in Vermont. Sometimes rose-color or nearly white.
- L. spicata*, Lam. In gravelly soil.
- L. Kalmii*, L. Near ponds and streams. Pomfret, *Morgan*; Sharon, *Dr. Baxter*; Chelsea and Olcott, Vt.; Olcott Falls, N. H., *Sargent*.
- L. inflata*, L. (Indian Tobacco.) Fields; very common.
- L. Dortmanna*, L. (Water Lobelia.) In ponds; New London, *Sargent*; Canaan, Plainfield, etc. Not very rare.

ORDER 44. CAMPANULACEÆ. CAMPANULA FAMILY.

Specularia, Heister. Venus's Looking-glass.

- S. perfoliata*, A. DC. Sterile grounds.

Campanula, Tourn. Bellflower.

- C. RAPUNCULOIDES*, L. Escaped from gardens; a European species.
- C. rotundifolia*, L. (Harebell.) Rocky or sandy banks; common.
- C. aparinoides*, Pursh. (Marsh Bellflower.) Wet meadows; in the grass.

ORDER 45. ERICACEÆ. HEATH FAMILY.

Gaylussacia, HBK.

- G. resinosa*, Torr. & Gray. (Huckleberry.) Swamps; Hanover and Lebanon, N. H.; Hartford, Vt.; not abundant.

Vaccinium, L. Blueberry. Bilberry. Cranberry.

- V. Pennsylvanicum*, Lam. (Low Blueberry.) Very common and ripe early.

- V. Canadense*, Kalm. (Canada Blueberry.) Swamps and low grounds; common.

- V. vacillans*, Solander. (Half-high Blueberry.) Hanover, N. H.; Hartford and Fairlee, Vt.; N. Pomfret, Vt., *Morgan*. Not common; intermediate in size between the preceding and the next.

- V. corymbosum*, L. (High or Swamp Blueberry.) Ripens later; abundant and variable.

- V. uliginosum*, L. (Bog Bilberry.) Moosilauke Mt., *Flint*.

- V. Vitis-Idæa*, L. (Cowberry. Mountain Cranberry.) Moosilauke Mt., *Flint*; Cardigan Mt., *A. Fairbanks*.

- V. Oxyccoccus*, L. (Small Cranberry.) Bogs; frequent.

- V. macrocarpon*, Ait. (Common Cranberry.) Bogs; common.

Chiogenes, Salisb. Creeping Snowberry.

- C. serpyllifolia*, Salisb. (*C. hispidula*, Torr. & Gray.) Bogs and mountain woods; abundant.

Arctostaphylos, Adans. Bearberry.

- A. Uva-ursi*, Spreng. Fairlee Cliffs, Vt., *F. O. Grover*; and not rare both north and south of our limits.

Epigæa, L. Trailing Arbutus. May-flower of New England.
E. repens, L. Woods and clearings; abundant.

Gaultheria, Kalm. Wintergreen or Checkerberry.
G. procumbens, L. Woods; very common.

Andromeda, L.

A. polifolia, L. Peat bogs. Lebanon, Meriden, Canaan, etc.
A. ligustrina, Muhl. Wet grounds; Enfield, *F. G. Atwell*; Sunapee Lake,
B. P. Ruggles.

Cassandra, Don. Leather-Leaf.

C. calyculata, Don. Swamps; common.

Kalmia, L. Laurel.

K. angustifolia, L. (Sheep Laurel.) Common.

K. glauca, Ait. (Pale Laurel.) Cold bogs. Lebanon, Meriden, Hanover, etc

K. latifolia, L. (Mountain Laurel. Spoon-wood.) Ashland, on Squam
 Lake, *Sargent*; more common southward.

Rhododendron, L. Rose Bay. Azalea. Rhodora.

R. nudiflorum, Torr. (Azalea nudiflora, L.)—(Purple Azalea.) Swamps;
 Lebanon and New London, N. H.; Norwich, Windsor, Ludlow, and
 Hartford, Vt.; not abundant.

R. viscosum, Torr. (Azalea viscosa, L.)—(White Azalea.) Sunapee Lake,
B. P. Ruggles. Not common.

R. Rhodora, Don. (Rhodora Canadensis, L.) Peat swamps. Meriden,
Dr. Barrows; Hanover, *Jesup*; Canaan, *Wallace*; Plymouth, *Sargent*.

R. maximum, L. (Great Laurel.) This superb plant has been found in
 North Grantham, *C. H. Hitchcock*, *Dr. Barrows*, *G. C. Barton*.

Ledum, L. Labrador Tea.

L. latifolium, Ait. Cold bogs and mountain woods. Lebanon, *E. A. Mack*; Bear Pond in Canaan, *Wallace*; Cardigan Mt., *A. Fairbanks*; New London, *Carter*; and probably not infrequent.

Loiseleuria, Desv. Alpine Azalea.

L. procumbens, Desv. Moosilauke Mt., *A. Fairbanks*.

Chimaphila, Pursh. Pipsissewa.

C. umbellata, Nutt. (Prince's Pine.) Dry woods.

C. maculata, Pursh. With the above.

Moneses, Salisb. One-flowered Pyrola.

M. grandiflora, Salisb. (*M. uniflora*, Gray.) Woods. Lebanon and Hanover, N. H.; Norwich and Royalton, Vt.

Pyrola, Tourn. Wintergreen. Shin-Leaf.

P. secunda, L. All the species common in rich woods.

P. chlorantha, Swartz. Woods.

P. elliptica, Nutt. (Shin-Leaf.) Woods.

P. rotundifolia, L. Woods; several varieties.

Pterospora, Nutt. Pine-drops.

P. Andromedea, Nutt. "Parasitic apparently on the roots of pines;" very rare. Lebanon, near Olcott Falls, *Jesup*.

Monotropa, L.

- M. uniflora, L. (Indian Pipe.) Dark woods; common.
 M. Hypopitys, L. (Pine-sap.) Under pine trees. North Pomfret, Vt.,
Morgan; Royalton, Vt., *Ward*.

ORDER 46. **PRIMULACEÆ. PRIMROSE FAMILY.****Trientalis, L.**

- T. Americana, Pursh. (Star-flower.) Wet woods.

Steironema, Raf.

- S. ciliatum, Raf. (Lysimachia ciliata, L.) Low grounds; common.
 S. lanceolatum, var. *hybridum*, Gray. (Lysimachia lanceolata, Walt.) Banks of Conn. river.

Lysimachia, Tourn. Loosestrife.

- L. quadrifolia, L. (Four-leaved Loosestrife.) Woods; common.
 L. stricta, Ait. (Strict Loosestrife.) Low grounds; common.
 L. thrysiflora, L. (Tufted Loosestrife.) Mud Pond, Plainfield, N. H.,
Dr. Barrows; Royalton, Vt., *Ward*.
 L. NUMMULARIA, L. (Moneywort.) A troublesome weed in dooryards;
 European.

Anagallis, Tourn. Pimpernel.

- A. ARVENSIS, L. An occasional weed in gardens; European.

ORDER 47. **OLEACEÆ. OLIVE FAMILY.****Fraxinus, Tourn. Ash.**

- F. Americana, L. (White Ash.) Common in rich woods.
 F. pubescens, Lam. (Red Ash.) Squam Lake, N. H.; Randolph, Vt.,
Bates; occasionally seen.
 F. sambucifolia, Lam. (Black or Brown Ash.) Swamps; common.

Ligustrum, Tourn. Privet.

- L. VULGARE, L. Naturalized in places. *Dr. Barrows*. From Europe;
 used for hedges.
 [CHIONANTHUS VIRGINICA (The Fringe-tree of the South), SYRINGA
 VULGARIS (Common Lilac), and S. PERSICA (Persian Lilac), are in common cultivation.]

ORDER 48. **APOCYNACEÆ. DOGBANE FAMILY.****Apocynum, Tourn.**

- A. androsæmifolium, L. (Dogbane.) Borders of thickets; common.
 A. cannabinum, L. (Indian Hemp.) River-banks; frequent.

ORDER 49. **ASCLEPIADACEÆ. MILKWEED FAMILY.****Asclepias, L. Milkweed. Silkweed.**

- A. tuberosa, L. (Butterfly-weed.) Dry fields.
 A. purpurascens, L. (Purple Milkweed.) Charlestown, N. H., *Dr. Graves*;
 Royalton, Vt., *Ward*.

- A. *incarnata*, L. (Swamp Milkweed.) Swamps; common.
var. *pulchra*, Pers. With the type.
- A. *Cornuti*, Decaisne. (Common Milkweed.) Everywhere.
- A. *obtusifolia*, Michx. Roadsides; not common.
- A. *phytolaccoides*, Pursh. (Poke Milkweed.) Common.
- A. *quadrifolia*, L. Woods; not rare.
- A. *verticillata*, L. *Geological Survey of N. H.*; probably rare.

Vincetoxicum, Moench.

V. *NIGRUM*, Moench. Windsor, Vt., *Leland*; rare, introduced from Europe.

ORDER 50. **GENTIANACEÆ**. GENTIAN FAMILY.**Gentiana**, Tourn. Gentian.

- G. *crinita*, Froel. (Fringed Gentian.) Moist meadows. Hanover Centre, Lyme, etc.; not abundant.
- G. *quinqueflora*, Lam. (Five-flowered Gentian.) Hillside woods and fields. Hanover, Lebanon, etc., *Jesup*; not very common.
- G. *Andrewsii*, Griseb. (Closed Gentian.) Wet ground; common.

Menyanthes, Tourn. Buckbean.

M. *trifoliata*, L. In ponds and bogs. Hanover and Meriden, N. H.; Norwich, Vt.; etc.

Limnanthemum, Gmelin. Floating-Heart.

L. *lacunosum*, Griseb. In ponds; common.

ORDER 51. **POLEMONIACEÆ**. POLEMONIUM FAMILY.**Phlox**, L. Phlox.

[P. *PANICULATA*, P. *REPTANS* and P. *SUBULATA* (Moss Pink), perennials; also P. *DRUMMONDI*, an annual;—all from the West and South; have long been cultivated in old gardens.]

Polemonium, Tourn. Greek Valerian.

[P. *CÆRULEUM* (Jacob's Ladder) of the Middle States, a rare native of New Hampshire, is often cultivated; as also is P. *REPTANS* (Greek Valerian) from the same region.]

ORDER 52. **HYDROPHYLLACEÆ**. WATERLEAF FAMILY.**Hydrophyllum**, Tourn. Waterleaf.

H. *Virginicum*, L. (Common Waterleaf.) Damp woods.

ORDER 53. **BORRAGINACEÆ**. BORAGE FAMILY.**Cynoglossum**, Tourn. Hound's-Tongue.

C. *OFFICINALE*, L. (Common Hound's-Tongue.) Pastures and roadsides; from Europe.

C. *Virginicum*, L. (Wild Comfrey.) Olcott Falls, etc.

Echinosperrum, Lehm. Stickseed.

E. *Virginicum*, Lehm. (*Cynoglossum Morisoni*, DC.)—(Beggar's Lice.) Borders of woods.

E. *LAPPULA*, Lehm. Waste heaps; European weed.

Myosotis, Dill. Forget-me-not.

M. laxa, Lehm. (*M. palustris*, var. *laxa*, Gray.) Hanover, N. H.; Sharon, Vt., *Dr. Baxter*; Randolph, Vt., *Rev. J. A. Bates*.

M. arvensis, Hoffm. Sharon, Vt., *Dr. Baxter*; Hanover, N. H.; a garden weed.

M. verna, Nutt. Windsor, Vt., *Leland*; not common.

[*M. PALUSTRIS*. Withering. This species, the true Forget-me-not of Europe, is cultivated, and possibly spontaneous.]

Lithospermum, Tourn. Gromwell.

L. ARVENSE, L. West Lebanon, *Dr. E. E. Phelps*; not often seen.

L. OFFICINALE, L. Roadsides and pastures; both species European weeds.

Symphytum, Tourn. Comfrey.

S. OFFICINALE, L. A not common garden escape; from Europe.

Echium, Tourn. Bugloss.

E. VULGARE, L. (Blue Thistle.) Hanover, N. H.; Pomfret, Sharon, and Royalton, Vt. Roadsides and fields; European.

Mertensia, Roth. Lungwort.

[*M. VIRGINICA*, DC. (Virginian Cowslip. Blue Bells.) A native of the West and often cultivated.]

ORDER 54. **CONVOLVULACEÆ**. CONVOLVULUS FAMILY.**Ipomoea**, L. Morning-Glory.

[*I. QUAMOCЛИT* (Cypress-Vine), *I. COCCINEA* (Scarlet Morning-Glory), and *I. PURPUREA* (Common Morning-Glory) are everywhere in gardens.]

Convolvulus, Tourn. Bindweed.

C. spithameus, L. (*Calystegia spithamea*, Pursh.) Dry fields; Hanover, N. H., *Jesup*; Norwich, Vt., *E. D. Kelley*.

C. sepium, L. (*Calystegia sepium*, R. Br.)—(Hedge Bindweed.) Banks of streams and variable.

var. *Americanus*, Sims, is glabrous and common.

var. *repens*, Gray, is pubescent.

C. ARVENTIS, L. A smaller procumbent plant and common in old fields.

[A double flowering species, sold by the florists as "CALYSTEGIA PUBESCENS from China," when once planted spreads rapidly.]

Cuscuta, Tourn. Dodder.

C. Gronovii, Willd. Low grounds; parasitic on herbs, etc.; not rare.

ORDER 55. **SOLANACEÆ**. NIGHTSHADE FAMILY.**Solanum**, Tourn. Nightshade.

S. DULCAMARA, L. (Bittersweet.) Naturalized from Europe.

S. nigrum, L. (Common Nightshade.) Shaded grounds.

Physalis, L. Ground Cherry.

P. Virginiana, Mill. (*P. viscosa*, Gray's Man., 5th ed.) Sandy soils.

P. pubescens, L. An occasional weed; neither species common.

Nicandra, Adans. Apple of Peru.

N. PHYSALOIDES, Gaertn. In waste ground, but not common. Native of Peru.

Lycium, L. Matrimony-Vine.

L. VULGARE, Dunal. Near dwellings where it was once cultivated; European.

Hyoscyamus, Tourn. Henbane.

H. NIGER, L. Rarely seen; roadsides, etc. From Europe.

Datura, L. Jamestown-weed. Jimson-weed.

D. STRAMONIUM, L. (Stramonium. Thorn Apple.) Waste heaps. From Asia.

D. TATULA, L. (Purple Thorn Apple.) Waste heaps. From tropical America.

[Cultivated species of this Order are:—*LYCOPERSICUM ESCULENTUM* (Tomato), *SOLANUM TUBEROSUM* (Common Potato), and *NICOTIANA TABACUM* (Tobacco).]

ORDER 56. SCROPHULARIACEÆ. FIGWORT FAMILY.

Verbascum, L. Mullein.

V. THAPSUS, L. (Common Mullein.) Fields everywhere.

V. BLATTARIA, L. (Moth Mullein.) Less common, both species European weeds.

Linaria, Tourn. Toad-Flax.

L. CANADENSIS, Dumont. Sandy soil; common.

L. VULGARIS, L. (Toad-Flax. Butter and Eggs.) Roadsides; European.

Scrophularia, Tourn. Figwort.

S. nodosa, L., var. *Mariolandica*, Gray. Roadsides; Lebanon, N. H., *Jesup*; Royalton, Vt., *Ward*.

Chelone, Tourn. Turtle-head. Snake-head.

C. glabra, L. Wet places; common.

Pentstemon, Mitchell. Beard-tongue.

P. pubescens, Solander. Windsor, Vt., *Leland*; Norwich, Vt., *G. A. Loveland*. Not common.

P. laevigatus, Solander, var. *Digitalis*, Gray. (*P. Digitalis*, Nutt.) Lebanon, perhaps introduced with grass-seed.

Mimulus, L. Monkey-flower.

M. ringens, L. Wet places; common.

Gratiola, L. Hedge-Hyssop.

G. Virginiana, L. Wet places.

G. aurea, Muhl. Margins of ponds. Meriden, *Dr. Barrows*.

Ilysianthes, Raf. False Pimpernel.

I. riparia, Raf. (*I. gratioloides*, Benth.) Sandy borders of ponds and streams.

Veronica, L. Speedwell.

- V. Virginica, L. (Culver's-root.) Roadsides in rich soil; not common, appearing like a garden escape.
- V. Anagallis, L. (Water Speedwell.) Ditches.
- V. Americana, Schwein. (American Brooklime.) Brooks and ditches.
- V. scutellata, L. (Marsh Speedwell.) Bogs.
- V. officinalis, L. (Common Speedwell.) Claremont, *Dr. Barrows*.
- V. serpyllifolia, L. (Thyme-leaved Speedwell.) With grass; common.
- V. peregrina, L. (Purslane Speedwell.) A weed in cultivated grounds.
- V. ARVENSIS, L. (Corn Speedwell.) Claremont, *Dr. Barrows*.
- V. BUxbaumii, Tenore. Hanover; a weed in gardens and lawns.

Gerardia, L. False Foxglove.

- G. pedicularia, L. Dry woods. Fairlee Cliffs, Vt., etc.
- G. flava, L. (Downy False Foxglove.) Dry woods.
- G. quercifolia, Pursh. (Oak-leaved Foxglove.) Wooded hillsides. Fairlee Cliffs, Vt., etc.
- G. tenuifolia, Vahl. (Slender Foxglove.) Dry woods. *Dr. Barrows*.

Pedicularis, Tourn. Lousewort.

- P. Canadensis, L. Hanover, etc. Woods and banks.

Melampyrum, Tourn. Cow-Wheat.

- M. Americanum, Michx. Woods. Common.

ORDER 57. **OROBANCHACEÆ.** BROOM-RAPE FAMILY.**Epiphegus,** Nutt. Beech-drops.

- E. Virginiana, Bart. Usually under Beech trees. Hanover and New London, N. H.; Norwich and Royalton, Vt., not rare.

Conopholis, Wallroth. Cancer-root.

- C. Americana, Wallroth. In old woods. Claremont, N. H., *Dr. Graves*; [West Rutland, Vt., *Denslow*, beyond our limits.]

Aphyllon, Mitchell. Naked Broom-rape.

- A. uniflorum, Gray. Borders of woods; not rare.

ORDER 58. **LENTIBULARIACEÆ.** BLADDERWORT FAMILY.**Utricularia,** L. Bladderwort.

- U. inflata, Walt. Claremont, *Dr. Graves*; ponds.
- U. vulgaris, L. (Common Bladderwort.) Ponds; common.
- U. minor, L. Claremont, *Dr. Barrows*.
- U. gibba, L. Claremont, *Dr. Barrows*; West Lebanon, *Dr. Kennedy*.
- U. intermedia, Hayne. Marshes and shallow pools; Canaan, etc.; not rare.
- U. purpurea, Walt. Goose Pond, Canaan; rare.
- U. cornuta, Michx. Sandy margins of ponds. Plainfield, Canaan, New London, etc.; rather common.

ORDER 59. VERBENACEÆ. VERVAIN FAMILY.

Verbena, Tourn. Vervain.

V. urticæfolia, L. (White Vervain.) Common.

V. angustifolia, Michx.. In an old pasture near Hanover; rare so far north.

V. hastata, L. (Blue Vervain.) Roadsides; common.

[The Verbenas of the gardens are hybrid forms derived from a number of South American species, including one (V. AUBLETIA) which is indigenous to the West and South.]

Phryma, L. Lopseed.

P. Leptostachya, L. Woods; common.

ORDER 60. LABIATÆ. MINT FAMILY.

Trichostema, L. Blue-curls. Bastard Pennyroyal.

T. dichotomum, L. Sandy fields.

Teucrium, Tourn. Germander.

T. Canadense, L. (Wood Sage.) Near Conn. river. Hanover, N. H.; Olcott, Vt.; etc.

Mentha, Tourn. Mint.

M. VIRIDIS, L. (Spearmint.) Wet places; common; European.

M. PIPERITA, L. (Peppermint.) Along brooks; frequent; European.

M. Canadensis, L. Like the above, in wet places; common.

M. SATIVA, L. (Whorled Mint.) Plymouth, *Sargent*. A garden escape, perhaps.**Lycopus**, Tourn. Water Horehound.

L. Virginicus, L. (Bugle-weed.) Marshes.

L. sinuatus, Ell. (L. Europæus, L., var. sinuatus, Gray.) Common in wet grounds.

Pyrenanthemum, Michx. Mountain Mint.P. lanceolatum, Pursh. Claremont, *Dr. Barrows*; not common.P. muticum, Pers. Royalton, Vt., *Hard*.P. incanum, Michx. Claremont, *Dr. Barrows*; rare.**Origanum**, Tourn. Wild Marjoram.O. VULGARE, L. Claremont, *Dr. Barrows*; rare; from Europe.**Calamintha**, Tourn. Calamint.

C. Clinopodium, Benth. (Wild Basil.) Low woods.

Hedeoma, Pers. American Pennyroyal.

H. pulegioides, Pers. Barren fields. Common.

Monarda, L. Horse-Mint.

M. didyma, L. In old gardens called " Bee Balm;" infrequent in a wild state.

M. punctata, L. (Horse-Mint.) Claremont, *Dr. Barrows*; introduced.M. fistulosa, L., var. mollis, Benth. (Wild Bergamot.) In gardens; indigenous in Vt., *Perkins*.

Blephilia, Raf.

B. hirsuta, Benth. Killington Mt., Vt., *Denslow*.

Nepeta, L.

N. CATARIA, L. (Catnip.) Near dwellings.

N. GLECHOMA, Benth. (Ground Ivy.) Dooryards, etc. Both species European weeds.

Scutellaria, L. Skullcap.

S. lateriflora, L. (Mad-dog Skullcap.) Wet shady places.

S. galericulata, L. With the preceding; both species common.

Brunella, Tourn. Self-heal.

B. vulgaris, L. Fields; common.

Physostegia, Benth. False Dragon-head.

P. Virginiana, Benth. Hartford, Vt.; Lebanon, N. H.; in gardens, sometimes escaping to the roadsides. Native in Vermont.

Marrubium, Tourn. Horehound.

M. VULGARE, L. Escaped from gardens into waste places; European.

Leonurus, L. Motherwort.

L. CARDIACA, L. Around dwellings; a weed from Europe.

Lamium, L. Dead-Nettle.

L. AMPLEXICAULE, L. In cultivated grounds; rare.

L. MACULATUM, L. A garden escape; both European.

Galeopsis, L. Hemp-Nettle.

G. TETRAHIT, L. (Common Hemp-Nettle.) Waste places; common.

G. LADANUM, L. (Red Hemp-Nettle.) Hanover, etc., *Flint*; rare; both European.

Stachys, Tourn. Hedge-Nettle.

S. palustris, L. Near streams.

S. aspera, Michx. (*S. palustris*, var. *aspera*, Gray.) Wet grounds; not uncommon.

[Old gardens furnish other Labiate plants, such as *OCIMUM BASILICUM* (Sweet Basil), *LAVANDULA VERA* (Lavender), *HYSSOPUS OFFICINALIS* (Hyssop), *THYMUS SERPYLLUM* (Creeping Thyme), *SATUREIA HORTENSIS* (Summer Savory), *MELISSA OFFICINALIS* (Balm), *SALVIA OFFICINALIS* (Common Sage), etc.;—all from Europe.]

ORDER 61. **PLANTAGINACEÆ.** PLANTAIN FAMILY.**Plantago**, Tourn. Plantain.

P. MAJOR, L. (Common Plantain.) A European weed.

P. Rugellii, Dec. (P. Kamtschatika, Gray's Man., 5th ed.) As common as the preceding and generally mistaken for it, *Jesup*. (Gray's Flora of N. A., Vol. II, Part I, p. 389.)

P. LANCEOLATA, L. (Rib-grass. English Plantain.) Dooryards and dry fields; common.

DIVISION III. APETALOUS EXOGENS.

ORDER 62. **ILLECEBRACEÆ.** KNOTWORT FAMILY.

(Formerly included in CARYOPHYLLACEÆ.)

Anychia, Michx. Forked Chickweed.A. *capillacea*, DC. (A. *dichotoma*, var. *capillacea*, Torr.) Dry woods.
*Geological Survey of N. H.***Scleranthus**, L. Knawel.S. *ANNUUS*, L. Waste places; from Europe.ORDER 63. **AMARANTACEÆ.** AMARANTH FAMILY.**Amarantus**, Tourn. Amaranth.

- A. *HYPOTHONDRIACUS*, L. (Red Amaranth.) Escaped from gardens.
 A. *PANICULATUS*, L. A weed in gardens.
 A. *RETROFLEXUS*, L. (Hogweed. Pigweed.) Cultivated ground.
 A. *CHLOROSTACHYS*, Willd. (A. *retroflexus*, var. *chlorostachys*, Gray.)
 Cultivated ground.
 var. *HYBRIDUS*, Watson. (A. *retroflexus*, var. *hybridus*, Gray.) All
 the above from tropical America.
 A. *albus*, L. (Tumble Weed.) A coarse weed.

[Cultivated species of this Order:—*CELOSIA CRISTATA* (Common Cocks-comb), and *GOMPHRENA GLOBOSA* (Globe Amaranth); both from India.]ORDER 64. **CHENOPODIACEÆ.** GOOSEFOOT FAMILY.**Chenopodium**, Tourn. Pigweed. Goosefoot.

- C. *POLYSPERMUM*, L. Claremont, *Dr. Barrows*. European.
 C. *ALBUM*, L. (Lamb's Quarters. Pigweed.) Very common. European.
 C. *URBICUM*, L. Claremont, *Dr. Barrows*. European.
 C. *hybridum*, L. (Maple-leaved Goosefoot.) Introduced from New York.
 C. *GLAUCUM*, L. (Oak-leaved Goosefoot.) European.
 C. *BONUS-HENRICUS*, L. (*Blitum Bonus-Henricus*, Reich.)—(Good King Henry.) European.
 C. *capitatum*, Watson. (*Blitum capitatum*, L.)—(Strawberry Blite.)
 Around dwellings; infrequent.
 C. *BOTRYS*, L. (Jerusalem Oak.) Claremont, *Dr. Barrows*; Hanover,
 etc.; European.
 C. *AMBROSIOIDES*, L. (Mexican Tea.) Claremont, *Dr. Barrows*; from
 tropical America.

Roubieva, Moquin.R. *MULTIFIDA*, Moq. (*Chenopodium multifidum*, L.) Claremont, *Dr. Barrows*. All the above plants of this Order are weeds in the vicinity of cultivated grounds or dwellings.[Cultivated species of this Order include:—*BETA VULGARIS* (Common Beet), and *SPINACHIA OLERACEA* (Common Spinach), both from Europe.]

ORDER 65. **PHYTOLACCACEÆ.** POKEWEEED FAMILY.**Phytolacca**, Tourn. Pokeweed.

P. decandra, L. Roadsides. *Dr. Barrows.* More common in southern New England.

ORDER 66. **POLYGONACEÆ.** BUCKWHEAT FAMILY.**Rumex**, L. Dock. Sorrel.

[R. PATIENTIA, L. (Patience Dock.) This plant of northern Vt., and central Mass., should be within our limits.]

R. Britanica, L. (R. orbiculatus, Gray)—(Great Water-Dock.) Swamps.

R. altissimus, Wood. (R. Britanica, Gray)—(Pale Dock.) Claremont, *Dr. Barrows.*

R. verticillatus, L. (Swamp Dock.) Killington Mt., Vt., *Denslow.*

R. CRISPUS, L. (Curled Dock.) Very common everywhere.

R. OBTUSIFOLIUS, L. (Bitter Dock.) Fields and roadsides.

R. SANGUINEUS, L. Bloody-veined Dock.) Royalton, Vt., etc., *Ward.*

R. ACETOSELLA, L. (Sheep Sorrel.) The last four are weeds in fields and pastures. European.

Rheum, L. Rhubarb.

R. RHAPONTICUM, L. (Garden Rhubarb or Pie Plant.) Escaped from cultivation. Native of Asia.

Polygonum, Tourn. Knotweed. Smartweed. Bindweed.

P. aviculare, L. (Knotgrass.) Roadsides, dooryards, etc. Common.

P. erectum, L. (P. aviculare, var. erectum, Roth.) With the above.

P. tenue, Michx. Dry fields.

P. Pennsylvanicum, L. Fields; very common.

P. amphibium, L. (Water Smartweed.) Windsor, Vt., *Leland*: Sharon and Hartford, Vt.; Hanover and Enfield, N. H.; etc. In or near ponds.

P. Muhlenbergii, Watson. (P. amphibium, var. terrestre, Gray's Man., 5th ed.) Border of ponds.

P. ORIENTALE, L. (Prince's Feather.) In old gardens; from India.

P. PERSICARIA, L. (Lady's Thumb.) Fields; European.

P. hydropiperoides, Michx. (Water-pepper.) Shallow water.

P. Hydropiper, L. (Water-pepper.)

P. acre, HBK. (Water Smartweed.)

P. Virginianum, L. Banks of Conn. river, Windsor, Vt., *Leland*.

P. arifolium, L. (Halberd-leaved Tear-thumb.) Wet grounds.

P. sagittatum, L. (Arrow-leaved Tear-thumb.) Wet grounds.

P. CONVOLVULUS, L. (Black Bindweed.) Fields; European.

P. ciliode, Michx. On fences, etc. Common.

P. dumetorum, L., var. scandens, Gray. (Climbing False Buckwheat.) Common.

[P. Hartwrightii, Gray. (Common in Vt., *Prof. Perkins.*) May belong here.]

Fagopyrum, Tourn. Buckwheat.

F. ESCULENTUM, Moench. (Buckwheat.) Old fields.

F. TATARICUM, Gaertn. (India Wheat.) Both species cultivated; Asiatic.

ORDER 67. ARISTOLOCHIACEÆ. BIRTHWORT FAMILY.

Asarum, Tourn. Wild Ginger.

A. Canadense, L. Rich woods.

Aristolochia, Tourn. Birthwort.

[A. SIPHO, L'Her. (Pipe Vine. Dutchman's Pipe.) A native of Penn. and Va., and sometimes planted for arbors.]

ORDER 68. LAURACEÆ. LAUREL FAMILY.

Sassafras, Nees.S. officinale, Nees. Plainfield, N. H., *Miss Stevens*; Springfield, Vt., *Rev. S. Hayward*; Charlestown, N. H.; Randolph, Orange, and Pomfret, Vt.; rare; southward it is common.**Lindera**, Thunb. Spice-wood. Benjamin-bush.[L. Benzoin, Blume. A shrub found in Vermont (*Prof. Perkins*) and probably within our limits.]

ORDER 69. THYMELÆACEÆ. MEZEREUM FAMILY.

Dirca, L. Leatherwood.

D. palustris, L. Rich woods; sometimes in swamps, common.

Daphne, L. Mezereum.

[D. MEZEREUM, L. An early blooming garden shrub from Europe.]

ORDER 70. LORANTHACEÆ. MISTLETOE FAMILY.

Arceuthobium, Bieb.A. pusillum, Peck. (Dwarf Mistletoe.) Found parasitic on *Picea nigra* (Black Spruce) in a swamp in Lebanon (1883); also near Bear Pond, Canaan (1883), *Jesup*. Probably rare. It is also found in the Adirondack region of N. Y., and is very injurious.

ORDER 71. SANTALACEÆ. SANDALWOOD FAMILY.

Comandra, Nutt. Bastard Toad-flax.C. umbellata, Nutt. Canaan, *J. B. Wallace*.

ORDER 72. EUPHORBIACEÆ. SPURGE FAMILY.

Euphorbia, L. Spurge.

E. maculata, L. A common weed in gardens.

E. Preslii, Guss. (E. hypericifolia, Gray's Man., 5th ed.) Roadsides and gardens.

E. CYPARISSIAS, L. (Cypress Spurge.) Roadsides. Often seen in cemeteries.

E. PEPLUS, L. Dooryards. Hanover, *Jesup*. The last two are European. [E. marginata, Pursh., from the West, is cultivated.]**Acalypha**, L. Three-seeded Mercury.

A. Virginica, L. A weed in cultivated ground; frequent.

Ricinus, L. Castor-oil Plant.

[*R. COMMUNIS*, L. Very common in cultivation.]

ORDER 73. **URTICACEÆ**. NETTLE FAMILY.**Ulmus**, L. Elm.

U. fulva, Michx. (Red Elm. Slippery Elm.) Banks of streams; not rare.

U. Americana, L. (Common Elm. American or White Elm.) Abundant.

U. racemosa, Thomas. (Corky White Elm.) Meriden, *Dr. Barrows*, where a few large trees are found, apparently self sown. It is sometimes planted, and at the West is a valuable tree and indigenous.

Celtis, Tourn. Hackberry. Nettle Tree. Sugarberry.

C. occidentalis, L. Banks of Conn. river throughout the district, but not abundant.

Cannabis, Tourn.

C. SATIVA, L. (Hemp.) Roadsides; escaped from cultivation; European.

Humulus, L. Hopvine.

H. Lupulus, L. Roadsides; frequent in cultivation.

Urtica, Tourn. Nettle.

U. gracilis, Ait. (Common Nettle.) Fence rows; common.

U. URENS, L. Claremont, N. H., *Dr. Barrows*; Royalton, Vt., *Ward*; scarce.

Laportea, Gaudichaud.

L. Canadensis, Gaudich. (Wood-Nettle.) Rich woods.

Pilea, Lindl.

P. pumila, Gray. (Clearweed.) In shady places near dwellings.

Bœhmeria, Jacq. False Nettle.

B. cylindrica, Willd. Shady and moist ground.

[*MORUS RUBRA*, L., the Red Mulberry tree, is a native of central New England. *MORUS NIGRA* from the Levant with black fruit, and the white-fruited *MORUS ALBA* from China are occasionally planted.]

ORDER 74. **PLATANACEÆ**. PLANE-TREE FAMILY.**Platanus**, L. Plane-Tree.

P. occidentalis, L. (Buttonwood or American Sycamore.) Meriden and Hanover, N. H.; Hartford, Hartland, and Sharon, Vt. Not abundant so far north, and commonly planted.

ORDER 75. **JUGLANDACEÆ**. WALNUT FAMILY.**Juglans**, L. Walnut.

J. cinerea, L. (Butternut.) Very common.

[*J. NIGRA*, L., the Black Walnut of the West, is often seen in private grounds.]

Carya, Nutt. Hickory.

- C. alba, Nutt. (Shell-bark Hickory.) Claremont, N. H., *Dr. Barrows*;
Royalton, Vt., *L. Wild.*
C. porcina, Nutt. (Pig-nut. Hog Walnut.) Common.
C. amara, Nutt. (Bitter-nut.) Claremont and Lebanon, *Dr. Barrows*.
"Along the Conn. river on both sides," *Flint.*

ORDER 76. **MYRICACEÆ.** SWEET-GALE FAMILY.**Myrica**, L.

- M. Gale, L. (Sweet Gale.) Swamps; Hanover, Sunapee Lake, etc.; not uncommon.
M. asplenifolia, Endl. (*Comptonia asplenifolia*, Ait.)—(Sweet Fern.) Dry hills; common.

ORDER 77. **CUPULIFERÆ.** OAK FAMILY.

(Including BETULACEÆ, THE BIRCH FAMILY.)

Betula, Tourn. Birch.

- B. lenta, L. (Black or Cherry Birch.) Throughout our limits, but less common north of Hanover.
B. lutea, Michx. f. (Yellow Birch.) Common.
B. populifolia, Ait. (B. alba, var. *populifolia*, Spach.)—(American White Birch. Gray Birch.) Not abundant, but very common in the centre of the state.
B. ALBA, L. (European White Birch.) Introduced at Hanover, where it spreads rapidly.
B. papyrifera, Marsh. (B. *papyracea*, Ait.)—(Paper or Canoe Birch. White Birch of northern New England.) Very common.

Alnus, Tourn. Alder.

- A. viridis, DC. (Green or Mountain Alder.) Banks of Conn. river, Hanover, etc.
A. incana, Willd. (Speckled Alder.) Common northward.
A. serrulata, Ait. (Smooth Alder.) Claremont, *Dr. Graves*. More abundant southward.

Corylus, Tourn. Hazelnut.

- C. rostrata, Ait. (Beaked Hazelnut.) Common.
C. Americana, Walt. (Wild Hazelnut.) Hartford, Vt., and Hanover, N. H.; not uncommon.

Ostrya, Micheli. Hop-Hornbeam.

- O. Virginica, Willd. (Lever-wood. Iron-wood.) Common.

Carpinus, L. Hornbeam. Iron-wood.

- C. Caroliniana, Walt. (C. Americana, Michx.)—(American Hornbeam. Blue or Water Beech.) Common.

Quercus, L. Oak.

- Q. alba, L. (White Oak.) More or less common on both sides of the Conn. river.

Q. coccinea, Wang., var. *tinctoria*, Gray. (Black Oak.) North Charles-town, *Flint*; and in a few towns in both states.

Q. rubra, L. (Red Oak.) The most abundant species. Other species on our borders may yet be found within our limits.

Castanea, Tourn. Chestnut.

C. sativa, Mill., var. *Americana*. (*C. vesca*, var. Gray's Man., 5th ed.) Infrequent. Claremont is nearly the northern limit; frequently planted and growing well in sheltered places.

Fagus, Tourn. Beech.

F. ferruginea, Ait. Varies greatly; wood light colored or reddish, as growing on dry or moist ground. Abundant.

ORDER 78. **SALICACEÆ. WILLOW FAMILY.**

Salix, Tourn. Willow.

S. nigra, Marsh. (Black Willow.) The only native species tree-like in size.

S. lucida, Muhl. (Shining Willow.) Wet places; very handsome shrub.

S. FRAGILIS, L. (Brittle Willow.) Large tree; from Europe; early introduced.

S. ALBA, L. (White Willow.) Both this and the above with many varieties, and hybrid forms.

S. BABYLONICA, Tourn. (Weeping Willow.) From Asia, and here known only in the pistillate form.

S. longifolia, Muhl. (Long-leaved Willow.) Banks of the Conn. river at Lebanon, N. H., and Windsor, Vt. Not very common.

S. rostrata, Richard. (*S. livida*, Wahl., var. *occidentalis*, Gray.)—(Beaked Willow.) Common.

S. discolor, Muhl. (Glaucous Willow.) Very common everywhere.

S. humilis, Marsh. (Prairie Willow.) Dry ground; common.

S. tristis, Ait. (Dwarf Gray Willow.) With the above.

S. sericea, Marsh. (Silky Willow.) Not rare.

S. petiolaris, Smith. With the preceding.

S. PURPUREA, L. (Purple Willow.) From Europe; used for basket-work.

S. cordata, Muhl. (Heart-leaved Willow.) Very common and variable.

[In addition to the above, other species may occur on our mountains and elsewhere. Numerous hybrid forms have been noted, making the precise determination of the species at times impracticable. *SALIX FRAGILIS* and *S. ALBA* are much intercrossed. *S. DISCOLOR* and *S. HUMILIS* hybridize, and small forms of the latter are not readily distinguished from *S. TRISTIS*. See description of Genus in Manual, 6th ed. by M. S. BEBB.]

Populus, Tourn. Poplar. Aspen.

[*P. ALBA*, L. (White Poplar. Abele Tree.) Cultivated from Europe, and in the pistillate form only.]

P. tremuloides, Michx. (American Aspen.) Everywhere common.

P. grandidentata, Michx. (Great-toothed Poplar.) Frequent.

P. balsamifera, L. (Balsam Poplar.) Abundant near streams.

var. *CANDICANS*, Gray. (Balm of Gilead.) Commonly cultivated; never spontaneous within this district, and perhaps European.

[*P. MONILIFERA*, Ait. (Cotton-wood.) In cultivation from the West; rarely spontaneous.]

[*P. DILATATA*, Ait. (Lombardy Poplar.) From Europe, in the pistillate form only, and freely spreading by suckers.]

ORDER 79. **EMPETRACEÆ. CROWBERRY FAMILY.**

Empetrum, Tourn. Crowberry.

E. nigrum, L. (Black Crowberry.) Moosilauke Mt., *Rev. J. Blake.*

ORDER 80. **CERATOPHYLLACEÆ. HORNWORT FAMILY.**

Ceratophyllum, L. Hornwort.

C. demersum, L. Goose Pond in Canaan, George Pond in Enfield, N. H.; Fairlee Pond, Vt.; etc. Not rare.

SUBCLASS II. GYMNOSPERMÆ.

ORDER SI. CONIFERÆ. PINE FAMILY.

Pinus, Tourn. Pine.

- P. *Strobos*, L. (White Pine.) Very common; the most valuable species.
 P. *rigida*, Miller. (Pitch Pine.) Sandy soil; more common farther south.
 P. *resinosa*, Ait. (Red or Norway Pine.) Sparingly found throughout our limits. "It is not a native of Europe, but was named from the town of Norway in Maine." *Robert Douglass*.
 P. *SYLVESTRIS*, L. (Scotch Pine. Scotch Fir.) Already spontaneous in Hanover.

Picea, Link. Spruce.

- P. *nigra*, Link. (*Abies nigra*, Poir.)—(Black Spruce.) Swamps and cold mountain woods; common.
 var. *rubra*, Engelm. With the above, "cones larger, bright red-brown." Better timber than the type, and called by some lumbermen "Yellow Spruce."
 P. *alba*, Link. (*Abies alba*, Michx.)—(White Spruce.) Newport, N. H., *Dr. Graves*; Windsor, Vt., *Leland*; Killington Mt., Vt., *Denslow*; and more or less throughout our limits.

Tsuga, Carrière. Hemlock.

- T. *Canadensis*, Carr. (*Abies Canadensis*, Michx.)—(Hemlock.) Common.

Abies, Link. Fir.

- A. *balsamea*, Miller. (Balsam Fir.) Damp woods and mountain swamps; common.

Larix, Tourn. Larch.

- L. *Americana*, Michx. (American Larch. Tamarack. Hackmatack.) Cold swamps; common.
 L. *EUROPEA*, L. (European Larch.) Widely planted, and in Hanover spreading freely.

Thuya, Tourn. Arbor Vitæ.

- T. *occidentalis*, L. Commonly called White Cedar. Common in swamps and on river-banks.

Juniperus, L. Juniper.

- J. *comunis*, L. (Common Juniper.) Claremont (*Dr. Barrows*), Hanover, Orford, Enfield, Sutton, etc., N. H.; Pomfret, Orange, Williams-town, Vt.; etc.
 J. *Virginiana*, L. (Red Cedar.) Hanover, N. H.; Fairlee, Vt., *F. O. Grover*; and in many places in the district, but not common.

Taxus, Tourn. Yew.

- T. *Canadensis*, Willd. (*T. baccata*, var. *Canadensis*, Willd.)—(Ground Hemlock.) Common.

[Among evergreens in cultivation will be found:—*PINUS AUSTRIACA* (Austrian Pine), *PICEA EXCELSA* (Norway Spruce), *ABIES PECTINATA* (European Silver Fir), etc.]

CLASS II. MONOCOTYLEDONOUS OR ENDOGENOUS
PLANTS.

ORDER 82. HYDROCHARIDACEÆ. FROG'S-BIT FAMILY.

Elodea, Michx. Water-weed.

E. Canadensis, Michx. (*Anacharis Canadensis*, Planchon.) Found in slow streams and ponds. In a pool near Conn. river, Norwich, Vt.; *Dr. Blanchard*; and doubtless elsewhere within our limits.

Vallisneria, L. Tape-Grass. Eel-Grass.

V. spiralis, L. Hanover, N. H., in Conn. river; Fairlee Pond, Vt.; etc.

ORDER 83. ORCHIDACEÆ. ORCHIS FAMILY.

Microstylis, Nutt. Adder's-Mouth.

M. monophyllos, Lindl. Hanover, N. H.; Pomfret, Union Village, and Ascutney Mt., Vt.

M. ophioglossoides, Nutt. Shrewsbury and Killington Mts., Vt., *Denslow*; Pomfret and Ascutney Mt., Vt.; New London, N. H., *Sargent*. Both species in wet woods.

Liparis, Richard. Twayblade.

L. liliifolia, Richard. Banks of the Conn. river at Windsor, Vt., *Leland*.
L. Læselii, Richard. Bogs and hillside pastures, Hanover, Meriden, Plymouth, etc.; not rare.

Calypso, Salisb. Calypso.

[*C. borealis*, Salisb. This rare and local plant of northern Vermont, one of the most beautiful of its Order, is said by *Dr. Blanchard* to be found in cold cedar swamps throughout Caledonia and Essex counties. He has himself collected it in Ryegate, Barnet, and Peacham, just beyond our northern limits, and it can hardly fail of being within these limits.]

Aplectrum, Nutt. Putty-root. Adam-and-Eve.

A. hiemale, Nutt. Walpole and Claremont, *Dr. Barrows*; New London, *W. F. Carter*. In rich, dark woods. Very rare.

Corallorrhiza, Haller. Coral-root.

C. innata, R. Br. Damp woods; not rare.

C. odontorhiza, Nutt. Norwich, Vt.; rare.

C. multiflora, Nutt. Rich woods. Hanover, etc., and on both sides of the Conn. river.

Listera, R. Brown. Twayblade.

L. cordata, R. Br. Moosilauke Mt., N. H., *Flint*; Killington Mt., Vt., *Denslow*.

L. convallarioides, Nutt. Mountain bogs.

Spiranthes, Richard. Ladies' Tresses.

S. latifolia, Torr. Moist banks. Windsor, Vt., *Leland*; Union Village, Vt., *Blanchard*; Norwich, Vt., *Sargent*; Royalton, Vt., *Ward*.

S. Romanzoffiana, Cham. North Pomfret, Vt., *Morgan*; Royalton, Vt., *Ward*; Grantham, N. H., *Barton*; New London, N. H., *Carter*.

S. cernua, Richard. Wet meadows; common.

S. gracilis, Big. Sandy soil. Hanover and Plymouth, N. H.; Norwich and Royalton, Vt.; etc.; not rare.

Goodyera, R. Br. Rattlesnake-Plantain.

G. repens, R. Br. In rich woods; not rare.

G. pubescens, R. Br. Rich woods; frequent.

Arethusa, Gronov. Arethusa.

A. bulbosa, L. Near Goose Pond in Canaan, N. H., *Dr. Barrows*; Peat bogs, Hanover, N. H., *Jesup*; Pine Hill Ponds, Rutland, Vt., *Denslow*: rare.

Calopogon, R. Br. Calopogon. Grass Pink.

C. pulchellus, R. Br. Bogs and moist pastures.

Pogonia, Juss. Pogonia.

P. ophioglossoides, Nutt. Bogs. Bear and Mud Ponds, Canaan; George Pond, Enfield; Hanover, etc., N. H.; Waterman's Pond near Union Village, Vt.; etc.

Orchis, L.

O. spectabilis, L. (Showy Orchis.) Rich woods; not rare.

[O. rotundifolia, Pursh. (*Habenaria rotundifolia*, Richard.) Sphagnum bogs, *Perkins's Vermont Flora*. This rare species is likely to occur on our northern borders.]

Habenaria, Willd. Rein-Orchis.

H. tridentata, Hook. Rich and moist woods; not rare.

H. virescens, Spreng. (Greenish Orchis.) Claremont, *Dr. Barrows*.

H. bracteata, R. Br. (H. viridis, var. bracteata, Reichenb.)—(Bracted Green Orchis.) Rich woods.

H. hyperborea, R. Br. (Northern Green Orchis.) Cold bogs, also wet woods.

H. dilatata, Gray. (Northern White Orchis.) Pomfret, Sharon, Norwich, Windsor, and Killington Mt., Vt.; Hanover, N. H.; etc. Wet grounds.

H. obtusata, Rich. Killington Mt., Vt., *Morgan*, *Denslow*.

H. Hookeri, Torr. (Smaller Round-leaved Orchis.) Norwich and Sharon, Vt., *Morgan*; Hanover and Lyme, N. H. Woods; not rare.

H. orbiculata, Torr. (Larger Round-leaved Orchis.) Mountain woods; not uncommon.

H. blephariglottis, Torr. (White Fringed Orchis.) Grantham Mt., and New London, N. H.; Royalton, Vt.; etc.

H. lacera, R. Br. Moist ground; common.

H. psycodes, Gray. (Purple Fringed Orchis.) Swamps; more common than the next.

H. fimbriata, R. Br. (Purple Fringed Orchis.) Claremont, *Dr. Barrows*; New London, *Sargent*. Spike not so dense, and flowers larger than the preceding, and blooming two weeks earlier.

Cypripedium, L. Lady's Slipper. Moccasin-Flower.

C. arietinum, R. Br. (Ram's-Head. Lady's Slipper.) Ascutney Mt., Vt.,
Dr. Phelps; Lyme, N. H., *R. S. Alden*; Hanover, N. H., near Olcott
 Falls, *E. P. Barrows*; very rare.

C. parviflorum, Salish. (Smaller Yellow Lady's Slipper.) Sometimes with
 the next, but not readily distinguished from it.

C. pubescens, Willd. (Larger Yellow Lady's Slipper.) Hanover, N. H.;
 Norwich, Vt.; frequent.

C. spectabile, Swartz. (Showy Lady's Slipper.) Swamps. Hanover, N.
 H., *F. H. Herrick*; Lebanon, N. H., *Dr. Barrows*; Sharon, Vt., *Dr.
 Baxter*; Barnard, Vt., *Ward*. Not very abundant.

C. acaule, Ait. (Stemless Lady's Slipper.) Dry or moist woods; com-
 mon.

ORDER 84. IRIDACEÆ. IRIS FAMILY.

Iris, Tourn. Blue-Flag.

I. versicolor, L. Swamps; common.

Sisyrinchium, L. Blue-eyed Grass.

S. angustifolium, Mill. (*S. Bermudiana*, var. *mucronatum*, Gray.) Moist
 meadows; common.

S. anceps, Cav. (*S. Bermudiana*, var. *anceps*, Gray.) Same range as the
 above.

[Garden plants of this Order belong to the following Genera:—**IRIS**,
TIGRIDIA (Tiger-flower), **GLADIOLUS** (Sword Lily), and **CROCUS**.]

ORDER 85. LILIACEÆ. LILY FAMILY.

Smilax, Tourn. Greenbrier. Cat-brier.

S. herbacea, L. (Carrion Flower.) River-banks; not infrequent.

S. rotundifolia, L. (Greenbrier.) Thickets; not common.

Allium, L. Onion.

A. tricoccum, Ait. (Wild Leek.) Rich woods; frequent on both sides of
 the Conn. river.

A. Schoenoprasum, L. (Chives.) Banks of the Conn. river at Windsor,
 Vt., *Leland*. In old gardens.

A. Canadense, Kalm. (Wild Garlic.) Moist meadows.

Ornithogalum, Tourn. Star-of-Bethlehem.

O. UMBELLATUM, L. Dooryards; common; European.

Hemerocallis, L. Day-lily.

H. FULVA, L. Roadsides, frequent and well established; from Europe.

Convallaria, L. Lily of the Valley.

[*C. MAJALIS*, L. Common in gardens; a native of Virginia, and some-
 times escaping to the roadsides. Our plant is of European origin.]

Polygonatum, Tourn. Solomon's Seal.

P. biflorum, Ell. (Common Solomon's Seal.) Wooded banks and hill-
 sides; common.

P. giganteum, Dietrich. (Great Solomon's Seal.) River-banks; rare; Hartford, Vt., *Ward*.

Asparagus, Tourn. Asparagus.

A. OFFICINALIS, L. A common garden escape; European.

Smilacina, Desf. False Solomon's Seal.

S. racemosa, Desf. (False Spikenard.) Moist copses.

S. stellata, Desf. Both banks of Conn. river; Plymouth, *Miss Langdon*; etc.

S. trifolia, Desf. Sphagnous swamps; somewhat rare.

Maianthemum, Wigg.

M. Canadense, Desf. (*Smilacina bifolia*, var. *Canadensis*, Gray.) Moist woods; very common.

Streptopus, Michx. Twist-Foot. Twisted-Stalk.

S. amplexifolius, DC. Claremont, Groton, Cube Mt., and Moosilauke Mt., N. H.; Pomfret and Killington Mt., Vt.; etc.

S. roseus, Michx. Frequent in damp woods.

Clintonia, Raf. Clintonia.

C. borealis, Raf. Moist woods; common.

Uvularia, L. Bellwort.

U. perfoliata, L. Claremont, *Dr. Barrows*; scarce.

U. grandiflora, Smith. Rich woods; not uncommon.

Oakesia, Watson. Sessile-leaved Bellwort.

O. sessilifolia, Watson. (*Uvularia sessilifolia*, L.) Common in low woods.

Erythronium, L. Adder's-Tongue.

E. Americanum, Ker. (Yellow Adder's-Tongue.) Very common in rich woodlands.

Lilium, L. Lily.

L. Philadelphicum, L. (Orange Lily.) Common in sandy, open woods.

L. Canadense, L. (Yellow Lily.) Moist meadows; very abundant.

[*L. TIGRINUM*, Ker. (Tiger Lily.) Sometimes spontaneous on the roadside. From Eastern Asia.]

Medeola, Gronov. Indian Cucumber-root.

M. Virginiana, L. Rich moist woods; common.

Trillium, L. Wake Robin. Birthroot.

T. erectum, L. (Early Trillium.) Common in rich woods.

var. *album*, Pursh. Hartford, Vt., *C. H. Hazen*; Chittenden, Vt., abundant, *Flint*. Found with the type.

T. grandiflorum, Salisb. (Large-flowered Trillium.) The finest species of the genus, and common in Vermont west of the Green Mts. It has been reported from Woodstock and Thetford, Vt., and should be looked for elsewhere on our western border.

T. cernuum, L. (Nodding Trillium.) Claremont, Hanover, and Lebanon, N. H.; Norwich, Vt., *H. A. Nichols*. Not abundant.

T. erythrocarpum, Michx. (Painted Trillium.) Moist woods; common.

Tofieldia, Hudson. False Asphodel.

T. glutinosa, Willd. Quechee Gulf, Vt., *Leland*; very rare.

Veratrum, Tourn. False Hellebore.

V. viride, Ait. (American White Hellebore. Indian Poke.) Low grounds; common.

[The following Liliaceous plants are found in gardens:—*COLCHICUM AUTUMNALE* from Europe, *LILIUM* of many species from Europe and the East, *FRITILLARIA IMPERIALIS* (Crown Imperial) from Asia, *TULIPA GESNERIANA* (Common Tulip of many varieties) also Asiatic, *SCILLA* (Squill) of several European species, *MUSCARI BOTRYOIDES* (Grape Hyacinth) from Europe, *HYACINTHUS ORIENTALIS* (Common Hyacinth) of the Levant, *FUNKIA* (Day Lily) from Europe and the East, *YUCCA FILAMENTOSA* of the south-western United States, etc.]

ORDER 86. **PONTEDERIACEÆ**. PICKEREL-WEED FAMILY.**Pontederia**, L. Pickerel-weed.

P. cordata, L. Ponds; common and trimorphous.

ORDER 87. **XYRIDACEÆ**. YELLOW-EYED-GRASS FAMILY.**Xyris**, Gronov. Yellow-eyed Grass.

X. flexuosa, Muhl. The typical form not yet reported.

var. *pusilla*, Gray. Sandy margins of streams and ponds. Meriden, Canaan, etc., *Jesup*.

ORDER 88. **JUNCACEÆ**. RUSH FAMILY.**Juncus**, Tourn. Rush.

J. effusus, L. Marshes. A large and very common species.

J. filiformis, L. Low grounds, not very rare.

J. trifidus, L. Moosilauke Mt., *Rev. J. Blake*.

J. marginatus, Rostk. Sands of Conn. river.

J. Greenii, O. & T. Haverhill, *Flint*.

J. tenuis, Willd. Very common.

J. bufonius, L. (Toad Rush.) Roadsides.

J. pelocarpus, E. Meyer. Shores of Enfield Pond, *Jesup*.

J. articulatus, L. Wet grounds.

J. acuminatus, Michx. Wet places; common and very variable.

J. nodosus, L. Swamps and banks of streams.

J. Canadensis, J. Gay. Common; a late blooming species.

var. *longicaudatus*, Engelm.

var. *coarctatus*, Engelm.

Luzula, DC. Wood-Rush.

L. vernalis, DC. (*L. pilosa*, Willd.) Hanover, *A. C. French*; Canaan, *Dr. Barrows*; not uncommon.

L. spadicea, DC., var. *melanocarpa*, Meyer. (*L. parviflora*, var. *melanocarpa*, Gray.) Moosilauke Mt., *Rev. J. Blake*.

L. campestris, DC. Meadows; common.

L. spicata, Desv. Moosilauke Mt., *Rev. J. Blake*.

ORDER 89. **TYPHACEÆ.** CAT-TAIL FAMILY.**Typha**, Tourn.

T. latifolia, L. (Common Cat-tail.)

Sparganium, Tourn. Bur-reed.

S. simplex, Huds. Borders of ponds and streams.

var. *androcladum*, Engelm. The largest form; common.

var. *fuitans*, Engelm. Not uncommon in ponds and streams.

var. *angustifolium*, Engelm. In ponds.

S. minimum, Fries. Pond holes, Norwich, Vt.; probably not rare.

ORDER 90. **ARACEÆ.** ARUM FAMILY.**Arisæma**, Martius.

A. triphyllum, Torr. (Indian Turnip.) Common in rich woods.

A. Dracontium, Schott. (Dragon Arum.) Charlestown, *Dr. Graves*; rare.

Peltandra, Raf. Arrow Arum.

P. undulata, Raf. (*P. Virginica*, Kunth.) Mud Pond, Canaan, *Jesup*: Enfield, *Atwell*. Perhaps not very common.

Calla, L. Water Arum.

C. palustris, L. Bogs. Claremont and Newport, *Dr. Barrows*; New London, N. H., *Carter*; Hartford, Springfield, and Sherburne, Vt.; not very common.

Symplocarpus, Salisb. Skunk Cabbage.

S. fætidus, Salisb. Wet places. Norwich and Thetford, Vt.; etc. Not as common as it is farther south.

Acorus, L. Sweet Flag. Calamus.

A. Calamus, L. Swamps; everywhere.

ORDER 91. **LEMNACEÆ.** DUCKWEED FAMILY.**Lemna**, L. Duckweed.

L. minor, L. Floating in still water; not infrequent in ponds and ditches. Hanover and Enfield, N. H.; Norwich, Vt.; etc.

Spirodela, Schleid. Duckweed.

S. polyrrhiza, Schleid. (*Lemna polyrrhiza*, L.) In the same localities as the preceding, and like it rarely ever found in flower or fruit; propagating by winter buds. This species is less common than the preceding.

ORDER 92. **ALISMACEÆ.** WATER-PLANTAIN FAMILY.**Alisma**, L. Water-Plantain.

A. Plantago, L. Shallow water; abundant and variable.

Sagittaria, L. Arrow-head.

S. variabilis, Engelm. In wet places; several varieties.

S. graminea, Michx. Lebanon, Hanover, etc.; not rare near ponds.

ORDER 93. **NAIADACEÆ.** PONDWEED FAMILY.**Scheuchzeria**, L.

S. palustris, L. Cranberry Pond on Grantham Mt., N. H.; Waterman's Pond near Union Village, Vt.; etc.; *Jesup*.

Potamogeton, Tourn. Pondweed.

P. natans, L. Ponds and streams.

P. Oakesianus, Robbins. Cranberry Pond on Grantham Mt.

P. Pennsylvanicus, Cham. (*P. Claytonii*, Tuckerm.) Common in ponds and streams.

P. Spirillus, Tuckerm. Streams and ponds.

P. hybridus, Michx. Shallow ponds.

P. rufescens, Schrad. Windsor, Vt.; infrequent.

P. fluitans, Roth. (*P. lonchites*, Tuck.) Conn. river; common.

P. amplifolius, Tuckerm. Ponds and large rivers.

P. heterophyllus, Schreb. (*P. gramineus*, Fries.) Conn. river; common and variable.

P. perfoliatus, L. Conn. river; common.

var. *lanceolatus*, Robbins. Conn. river.

P. zosteræfolius, Schum. (*P. compressus*, Fries.) Randolph, Vt., *J. W. Freley*; Fairlee Pond, Vt.; George Pond, Enfield, N. H.

P. pauciflorus, Pursh. Pools.

P. pusillus, L. Shallow pools; not rare.

P. Tuckermani, Robbins. Claremont, *Dr. Barrows*; scarce.

P. pectinatus, L. In Conn. river; abundant.

P. Robbinsii, Oakes. Mascoma Creek near West Lebanon, and George Pond, Enfield, N. H.; Fairlee Pond, Vt. Common.

[Other species than the above are probably within our limits.]

Zannichellia, Micheli. Horned Pondweed.

Z. palustris, L. Ponds. Claremont, *Dr. Barrows*; scarce.

Naias, L. Naiad.

N. flexilis, Rostk. & Schmidt. Ponds; rather common.

ORDER 94. **ERIOCAULEÆ.** PIPEWORT FAMILY.**Ericaulon**, L. Pipewort.

E. septangulare, With. Sandy borders of ponds; common.

ORDER 95. **CYPERACEÆ.** SEDGE FAMILY.**Cyperus**, Tourn. Galingale.

C. diandrus, Torr. Low ground, and variable.

C. aristatus, Rottb. (*C. inflexus*, Muhl.) Shores of Conn. river.

C. filiculmis, Vahl. Dry, sterile soil; common.

C. dentatus, Torr. Margins of ponds.

C. esculentus, L. (*C. phymatodes*, Muhl.) Low ground, spreading by root-like tubers.

C. strigosus, L. Very common.

Dulichium, Pers.

D. spathaceum, Pers. Marshes; common.

Eleocharis, R. Br. Spike-Rush.

E. Robbinsii, Oakes. In ponds, rather rare; George Pond, Enfield; etc.

E. ovata, R. Br. (*E. obtusa*, Schultes.) Wet places.

E. olivacea, Torr. Mouth of Mascoma river, Lebanon, *Jesup*.

E. palustris, R. Br. Wet places, and very variable in size, etc.

E. intermedia, Schultes. N. Pomfret, Vt., *A. P. Morgan*.

E. tenuis, Schultes. Wet places; very common.

E. acicularis, R. Br. Muddy shores; abundant.

Fimbristylis, Vahl.

F. autumnalis, Roem. & Schultes. Low grounds.

F. capillaris, Gray. Sandy fields.

Scirpus, Tourn. Bulrush or Club-Rush.

S. pungens, Vahl. Borders of ponds; common.

S. Torreyi, Olney. Borders of ponds.

S. lacustris, L. (*S. validus*, Vahl.)—(Great Bulrush.) Very common.

S. debilis, Pursh. Near ponds throughout the district.

S. sylvaticus, L., var. *digynus*, Boeckl. (*S. microcarpus*, Presl.) Along brooks; common.

S. atrovirens, Muhl. Wet meadows; common.

Eriophorum, L. Cotton-Grass.

E. cyperinum, L. (*Scirpus Eriophorum*, Michx.) Wet meadows; quite variable; lateral spikelets sometimes long pedicelled.

E. alpinum, L. Cold bogs. Not rare; an early species.

E. vaginatum, L. Peat bogs, Lebanon; not very common.

E. Virginicum, L. Low meadows; a variety has white wool.

E. polystachyon, L. Low meadows. Somewhat variable; very common.

E. gracile, Koch. Low meadows; frequent.

Rhynchospora, Vahl. Beak-Rush.

R. fusca, Roem. & Schultes. Cranberry Pond on Grantham Mt.; rare.

R. alba, Vahl. Bogs; common.

R. glomerata, Vahl. Low grounds; common.

Cladium, P. Browne. Twig-Rush.

C. mariscoides, Torr. Common in bogs.

Carex, Ruppius. Sedge.

C. aquatilis, Wahl. (35)¹ Low grounds, *Dr. Barrows*.

C. arctata, Boott. (51) Woods; common.

C. aurea, Nutt. (79) Moist banks; not rare.

C. bromoides, Schk. (124) Bogs; common.

C. bullata, Schk. (14) Canaan, *Dr. Barrows*; rare.

C. canescens, L. (119) Swamps; common.

¹ The number following each name refers to its position in Gray's Manual of Botany, 6th edition.

- var. *vulgaris*, Bailey.
var. *alpicola*, Wahl. (*C. vitilis*, Fries.)
C. castanea, Wahl. (49)—(*C. flexilis*, Rudge.) Pomfret, Vt., *Morgan*.
C. cephalophora, Muhl. (114) Fields; very abundant.
C. communis, Bailey. (89)—(*C. varia* of Man., 5th ed.) Hillsides; common.
C. conoidea, Schk. (65) Moist meadows.
C. crinita, Lam. (41)—(*C. gynandra*, Schw.) Swamps; variable.
C. debilis, Michx., var. *Rudgei*, Bailey. (52)—(*C. debilis*, Man., 5th ed.) Somewhat variable; common.
C. Deweyana, Schw. (123) Woods and clearings; frequent.
C. digitalis, Willd. (69) Hillsides.
C. echinata, Murray, var. *microstachys*, Boeckl. (118)—(*C. scirpoides*, Schk. and *C. sterilis*, Willd.) Very common and variable.
C. filiformis, L. (27) Peat bogs; frequent.
var. *latifolia*, Boeckl. (*C. lanuginosa*, Michx.) Low meadows; common.
C. flava, L. (62) Wet meadows; common.
var. *viridula*, Bailey. (*C. Ederi*, Ehrh.) Vermont:—banks of White River, *Flint*; Fairlee Lake, *Blanchard*; Pomfret, *Morgan*.
C. foenea, Willd. (130)—(*C. adusta*, Boott.) In clearings. Claremont, etc., *Jesup*.
C. folliculata, L. (4) Bogs; frequent.
C. formosa, Dewey. (56) Claremont, *Dr. Barrows*.
C. gracillima, Schw. (55) Wet meadows; abundant.
C. granularis, Muhl. (60) Lebanon, N. H., *Dr. Barrows*; in Vermont also, *Dr. Blanchard*.
C. grisea, Wahl. (58) Moist grounds.
C. Hitchcockiana, Dewey. (67) Pomfret, Vt., *A. R. Morse*: rare.
C. Houghtonii, Torr. (26) Norwich, Vt., *E. A. Edmands*; rare.
C. hystricina, Muhl. (18) Wet meadows; frequent.
C. intumescens, Rudge. (5) Swamps; very common.
C. laxiculmis, Schw. (70)—(*C. retrocurva*, Dewey.) Woods.
C. laxiflora, Lam. (68) Woods; several varieties; common.
C. lenticularis, Michx. (36) Shores of Enfield Pond, *Dr. Kennedy*; Baker's Pond, Orford, *Grover*.
C. limosa, L. (45) Peat bogs; not uncommon.
C. longirostris, Torr. (48) Lebanon, *H. C. Fall*. Frequent.
C. lupulina, Muhl. (7) Wet grounds; rather common, and variable.
C. lurida, Wahl. (16)—(*C. tentaculata*, Muhl.) Meadows; very common, and variable; hybridizes with the preceding.
C. Magellanica, Lam. (43)—(*C. irrigua*, Smith.) Norwich, Vt.; bogs.
C. monile, Tuckerm. (12) Moist grounds; frequent.
C. Muhlenbergii, Schk. (112) Sterile soil.
C. pallescens, L. (63) Meadows; common.
C. pauciflora, Lightf. (1) Sphagnous bogs. Plainfield, *Dr. Barrows*. Rare.

- C. pedunculata, Muhl. (82) Dry woods; common.
 C. Pennsylvanica, Lam. (88) Woods and hillsides; very common, and blooming early.
 C. plantaginea, Lam. (74) Rich hillside woods; common.
 C. platyphylla, Carey. (72) Rich hillside woods; common.
 C. polytrichoides, Muhl. (96) Swamps; abundant.
 C. prasina, Wahl. (39)—(C. miliacea, Muhl.) Bogs; not common. Borders of streams, *Dr. Barrows*.
 C. Pseudo-Cyperus, L. (19) Goose Pond in Canaan, *Dr. Barrows*.
 var. Americana, Hochst. (C. comosa, Boott.) Wet places.
 C. pubescens, Muhl. (92) Lebanon, *Dr. Barrows*.
 C. retrorsa, Schw. (15) Marshes, near streams.
 var. Hartii, Gray. Frequent.
 C. riparia, Curtis. (29) Borders of ponds, etc.; common.
 C. rosea, Schk. (110) Woods and meadows; very variable.
 C. scabrida, Schw. (23) Swamps; common.
 C. scoparia, Schk. (128) Meadows; common and variable.
 C. sparganioides, Muhl. (111) Low grounds; frequent.
 C. stipata, Muhl. (101) Low grounds; common.
 C. straminea, Willd. (132)—(C. straminea, var. tenera, 5th ed. of Man.)
 Copse and fields; extremely variable. (See Manual.)
 C. stricta, Lam. (34) Wet meadows; very abundant.
 var. decora, Bailey. (C. aperta, Boott.) Not as common as the type.
 C. tenella, Schk. (109) Cold swamps; common.
 C. teretiuscula, Gooden. (104) Norwich, Vt.; swamps.
 C. torta, Boott. (37) Banks of streams; not rare.
 C. tribuloides, Wahr. (127) (C. lagopodioides, Schk.) Wet grounds;
 many varieties.
 var. cristata, Bailey. (C. cristata, Schw.) Moist grounds.
 C. trisperma, Dewey. (122) Cold bogs; common.
 C. Tuckermani, Dewey. (13) Swamps; not uncommon.
 C. umbellata, Schk. (90) Rocky hillsides.
 C. utricularia, Boott. (11) Swamps; common, and variable.
 C. varia, Muhl. (86)—(C. Emmonsii, Dew.) Woods and hillsides.
 var. colorata, Bailey. With the type.
 C. virescens, Muhl. (46) Rocky woods.
 C. vulgaris, Fries. (33) Claremont, *Dr. Barrows*.
 C. vulpinoidea, Michx. (107) Low meadows; abundant.

ORDER 96. GRAMINEÆ. GRASS FAMILY.

Paspalum, L.

P. setaceum, Michx. Sandy fields.

Panicum, L. Panic-Grass.

P. GLABRUM, Gaudin. (Smooth Crab-Grass.) Gravelly soil. This and the next are European weeds.

P. SANGUINALE, L. (Crab-Grass. Finger-Grass.) Too common.

- P. capillare*, L. (Old-witch Grass.) In dry soil, everywhere.
P. virgatum, L. (Tall Panic-Grass.) River-banks, etc.; frequent.
P. xanthophyllum, Gray. (Yellow Panic-Grass.) Claremont, *Dr. Barrows*; Hanover and Lebanon; rare.
P. latifolium, L. (Broad-leaved Panic-Grass.) Low thickets; frequent.
P. clandestinum, L. Moist thickets; common.
P. depauperatum, Muhl. Dry woods; frequent.
P. dichotomum, L. Many varieties and abundant.
P. laxiflorum, Lam. Resembles the preceding, and found with it.
P. CRUS-GALLI, L. (Barnyard-Grass.) On waste heaps and in rich soil; very variable as to size and hairiness. From Europe.

Setaria, Beauv. Bristly Foxtail Grass.

- S. VERTICILLATA*, Beauv. Claremont, *Dr. Barrows*; not common. All the species are European.
S. GLAUCIA, Beauv. (Foxtail.) Cultivated grounds; common.
S. VIRIDIS, Beauv. (Green Foxtail.) With the preceding.
S. ITALICA, Kunth. (Millet. Hungarian Grass.) Rarely spontaneous.

Cenchrus, L. Bur-Grass. Hedgehog-Grass.

- C. tribuloides*, L. Sandy soil; the burs prickly, with sharp barbed spines.

Leersia, Swartz. White Grass.

- L. Virginica*, Willd. (White Grass.) Wet woods; frequent.
L. oryzoides, Swartz. (Rice Cut-Grass.) Marshes, and common.

Andropogon, Royen. Beard-Grass.

- A. furcatus*, Muhl. (Blue stem.) Dry and sterile soil; common.
A. scoparius, Michx. (Broom-grass.) Like the last, and with the same range.

Chrysopogon, Trin.

- C. nutans*, Benth. (Sorghum nutans, Gray.)—(Indian Grass. Wood-Grass.) Dry soil; with the preceding species.

Phalaris, L. Canary-Grass.

- P. arundinacea*, L. (Reed Canary-Grass.) Wet grounds; common.
var. *PICTA*, from Europe, is the Ribbon-Grass of gardens.

Anthoxanthum, L.

- A. ODORATUM*, L. (Sweet Vernal-Grass.) Meadows; from Europe.

Hierochloe, Gmelin. Holy-Grass.

- H. borealis*, Roem. & Schultes. (Vanilla or Seneca Grass.) Hanover, etc., *Flint*; rare.

Aristida, L. Triple-awned Grass.

- A. dichotoma*, Michx. (Poverty-Grass.) Sandy soil.

Oryzopsis, Michx. Mountain Rice.

- O. melanocarpa*, Muhl. (Black Mountain Rice.) Rocky woods; blooming in May; common.

- O. asperifolia*, Michx. (White Mountain Rice.) This and the next less common than the first species.

O. Canadensis, Torr. (Canadian Mountain Rice.) Hills and dry plains.

Milium, Tourn.

M. effusum, L. (Millet-Grass.) Damp woods; frequent.

Muhlenbergia, Schreber. Drop-seed Grass.

M. sobolifera, Trin. Rocky woods; frequent.

M. glomerata, Trin. (Spiked Muhlenbergia.) Wet banks; not rare.

M. Mexicana, Trin. (Mexican Muhlenbergia.) Common everywhere.

M. sylvatica, Torr. & Gray. (Wood Muhlenbergia.) Rocky woods; rather scarce.

Brachyelytrum, Beauv.

B. aristatum, Beauv. Rocky woods; very common.

Phleum, L. Cat's-tail Grass.

P. PRATENSE, L. (Timothy. Herd's-Grass in New England.) A most valuable European grass.

Alopecurus, L. Foxtail Grass.

A. PRATENSIS, L. (Meadow Foxtail.) Claremont, *Dr. Barrows*; moist meadows.

A. GENICULATUS, L. (Floating Foxtail.) Claremont, *Dr. Barrows*. This and the above, European.

var. *aristulatus*, Torr. (*A. aristulatus*, Michx.) A native form found in wet places. Neither the species nor the variety is common.

Sporobolus, R. Br. Drop-seed Grass. Rush-Grass.

S. vaginæflorus, Vasey. (*Vilfa vaginæflora*, Torr.) Barren fields.

S. serotinus, Gray. Wet places; common.

Agrostis, L. Bent-Grass.

A. ALBA, L. (White Bent-Grass.) Cultivated from Europe.

var. *VULGARIS*, Thurb. (Common Red-top. Herd's-Grass of Penn.) Naturalized from Europe. The type and the variety are both good grasses, but cannot always be distinguished.

A. perennans, Tuckerm. (Thin-Grass.) In damp shade.

A. scabra, Willd. (Hair-Grass.) Dry places; common.

A. canina, L. (Brown Bent-Grass.) Meadows.

Cinna, L. Wood Reed-Grass.

C. arundinacea, L. Moist woods; frequent.

C. pendula, Trin. (*C. arundinacea*, var. *pendula*, Gray.) Wet rocks.

Calamagrostis, Adans. Reed Bent-Grass.

C. Canadensis, Beauv. (*Deyeuxia Canadensis*, Hook.)—(Blue-Joint Grass.) Wet grounds; a good grass.

C. Nuttalliana, Steud. (*Deyeuxia Nuttalliana*, Vasey.) Windsor, Vt., *Leland*; rare.

C. Langsdorffii, Trin. (*Deyeuxia Langsdorffii*, Kunth.) Moosilauke Mt.

Arrhenatherum, Beauv. Oat-Grass.

A. AVENACEUM, Beauv. Windsor, Vt., *Leland*; European.

Holcus, L. Meadow Soft-Grass.

H. LANATUS, L. (Velvet-Grass.) European.

Deschampsia, Beauv. Hair-Grass.

- D. flexuosa, Trin. (*Aira flexuosa*, L.)—(Common Hair-Grass.) Dry places.
 D. cæspitosa, Beauv. (*Aira cæspitosa*, L.)—(Tufted Hair-Grass.) Shores of Conn. river, etc.

Avena, Tourn. Oat-Grass.

- A. striata, Michx. Rocky hillsides; frequent.
 [A. SATIVA, L. (Cultivated Oat.)]

Danthonia, DC. Oat-Grass. White-Grass.

- D. spicata, Beauv. Dry and poor soil; hillside pastures, etc.; common.
 Sometimes called "June Grass," a name that belongs to *Poa pratensis* (Kentucky Blue-Grass).

Eatonia, Raf.

- E. Pennsylvanica, Gray. Moist woods and rocks; frequent.

Eragrostis, Beauv.

- E. capillaris, Nees. Sandy soil; common.
 E. pectinacea, Gray. Dry, sandy fields; rare.

Dactylis, L. Orchard-Grass.

- D. GLOMERATA, L. Very common; European.

Poa, L. Meadow-Grass. Spear-Grass.

- P. ANNUA, L. (Low Spear-Grass.) Cultivated grounds; everywhere; naturalized from Europe.
 P. COMPRESSA, L. (Wire-Grass.) Dry, sterile soil; European and very common.
 P. nemoralis, L. (*P. cæsia*, Smith.) Thetford, Vt., Dr. Blanchard; rare.
 P. serotina, Ehrh. (Fowl Meadow-Grass.) Wet meadows, and valuable.
 P. pratensis, L. (June Grass. Kentucky Blue-Grass.) Common, and of value.
 P. TRIVIALIS, L. (Roughish Meadow-Grass.) In wet meadows; European, and not very common.
 P. alsodes, Gray. Hillside woods, Dr. Barrows. Rare.

Glyceria, R. Br. Manna-Grass.

- G. Canadensis, Trin. (Rattlesnake-Grass.) Wet places; very common.
 G. elongata, Trin. Wet woods; rare.
 G. nervata, Trin. Wet places; common.
 G. pallida, Trin. Shallow water.
 G. grandis, Watson. (*G. aquatica* of Am. authors; *G. arundinacea*, Kth.)—(Reed Meadow-Grass.) Common in marshes.
 G. fluitans, R. Br. Floating in shallow water.

Festuca, L. Fescue-Grass.

- F. tenella, Willd. Sterile soil; not common.
 F. ovina, L. (Sheep's Fescue.) Forms tufts in meadows, and varies greatly.
 F. nutans, Willd. (Nodding Fescue.) Frequent in rocky woods.
 F. ELATIOR, L. (Meadow Fescue.) A very useful grass; common in cultivation; from Europe.

Bromus, L. Brome-Grass.

- B. Kalmii, Gray. (Wild Chess.) Borders of woodlands.
- B. **SECALINUS**, L. (Chess or Cheat.) Grain fields; a pernicious European weed, but not very common.
- B. ciliatus, L. Borders of woods; common and variable.

Lolium, L. Darnel.

- L. **PERENNE**, L. (Common Darnel. Italian Rye-Grass.) Fields; a valuable European grass.

[*LOLIUM TEMULENTUM*, L. (Poison Darnel), also European, whose seeds are reputed to be poisonous, may yet be found in grain fields.]

Agropyrum, Gaertn.

- A. repens, Beauv. (*Triticum repens*. L.)—(Couch or Witch-Grass.) Everywhere common and variable.

Elymus, L. Wild Rye.

- E. Virginicus, L. River-banks; common.
- E. Canadensis, L. River-banks; common.
- E. striatus, Willd. Rocky woods; and like the above somewhat variable

Asprella, Willd. Bottle-brush Grass.

- A. *Hystrix*, Willd. (*Gymnostichum Hystrix*, Schreb.) Woods; common.

[There belong also to this Order, *TRITICUM VULGARE* (Wheat), *SECALE CEREALE* (Rye), *HORDEUM VULGARE* (Barley), and *ZEA MAYS* (Maize, Indian Corn), each including many varieties. In addition, many foreign Grasses are cultivated for ornamental purposes.]

SERIES 2.

CRYPTOGAMOUS OR FLOWERLESS PLANTS.

CLASS III. ACROGENS.

ORDER 97. **EQUISETACEÆ.** HORSETAIL FAMILY.

- Equisetum**, L. Horsetail. Scouring Rush.
E. arvense, L. (Common Horsetail.) Moist, especially gravelly soil; very common.
E. sylvaticum, L. (Wood Horsetail.) Wet shady places; common.
E. limosum, L. (Marsh Horsetail.) In shallow water; common.
E. hyemale, L. (Scouring Rush.) Wet, sandy banks; very common.
E. variegatum, Schl. Sandy banks of rivers. Hanover, N. H.; Hartford, Pomfret, and Union Village, Vt.; rare.
E. scirpoideus, Michx. Wooded hillsides; common.

ORDER 98. **FILICES.** FERNS.

- Polypodium**, L. Polypody.
P. vulgare, L. On rocks in woods; common.
Adiantum, L. Maidenhair Fern.
A. pedatum, L. Rich, moist woods; common.
Pteris, L. Brake.
P. aquilina, L. (Common Brake.) Pastures and woods; everywhere.
Pellæa, Link. Cliff-Brake.
P. gracilis, Hook. Wet, limestone rocks. Norwich, Vt., *Dr. Edward Hyde*; rare.
Asplenium, L. Spleenwort.
A. Trichomanes, L. Shaded rocks; common.
A. ebeneum, Ait. Rocky woods; common.
A. angustifolium, Michx. Rich, moist woods. Hanover, N. H., *Miss Hitchcock*; Windsor and Pomfret, Vt. Not very abundant.
A. thelypteroides, Michx. Rich woods; common.
A. Filix-fœmina, Bernh. (Lady-Fern.) Frequent and variable.

Camptosorus, Link. Walking-Leaf.

C. rhizophyllus, Link. On shaded rocks; very local. Lebanon, N. H.,
Dr. Barrows; Windsor, Vt., *Leland*; Norwich, Vt., *Miss Loveland*.

Phegopteris, Féé. Beech Fern.

P. polypodioides, Féé. Damp woods; common.

P. hexagonoptera, Féé. Norwich, Vt., *F. H. Herrick*; scarce.

P. Dryopteris, Féé. Rocky woods; common.

Aspidium, Swartz. Shield Fern.

A. Thelypteris, Swartz. Marshes; abundant.

A. Noveboracense, Swartz. Wet woods; abundant.

A. spinulosum, Swartz. Damp woods. This is the European type, and is rare.

var. *intermedium*, Gray. Common.

var. *dilatatum*, Gray. Summit of Mt. Ascutney, Vt., *Leland*; Killington Mt., *Denslow*.

A. Boottii, Tuckerm. (*A. spinulosum*, var., *Boottii*, Man., 5th ed.) Not rare in damp woods.

A. cristatum, Swartz. Swamps; common.

var. *Clintonianum*, D. C. Eaton. Hanover, N. H., *Miss Hitchcock*; and probably not very rare.

A. Goldianum, Hook. Rich and moist woods; Hanover, N. H., *Miss Hitchcock*; not abundant.

A. marginale, Swartz. Rocky hillsides, etc.; abundant.

A. acrostichoides, Swartz. (Christmas Fern.) Woods; common and variable.

A. aculeatum, Swartz., var. *Braunii*, Koch. North Pomfret and Killington Mt., Vt., *Morgan*. "The Gulf" in Weathersfield, Vt., *Leland*; rare.

Cystopteris, Bernh. Bladder Fern.

C. fragilis, Bernh. Shaded ravines; very common.

C. bulbifera, Bernh. Moist cliffs and rich woods; frequent.

Onoclea, L.

O. sensibilis, L. (Sensitive Fern.) Very common.

O. Struthiopteris, Hoff. (*Struthiopteris Germanica*, Willd.)—(Ostrich Fern.) Rich alluvial soil; abundant.

Woodsia, R. Br.

W. Ilvensis, R. Br. On rocks; common.

W. obtusa, Torrey. Rocky banks; rather rare.

Dicksonia, L'Her.

D. pilosiuscula, Willd. (*D. punctilobula*, Kunze.) Common in moist, shady places and pastures.

Osmunda, L. Flowering Fern.

O. regalis, L. (Queen-Fern. Royal-Fern.) Wet woods; rather common.

O. Claytoniana, L. (Clayton's Fern.) Wet woods; very common.

O. cinnamomea, L. (Cinnamon Fern.) Wet woods; common.

ORDER 99. **OPHIOGLOSSACEÆ.** ADDER'S-TONGUE FAMILY.

Botrychium, Swartz. Grape-Fern. Moonwort.

B. simplex, Hitch. Hillside pastures; very seldom met with.

B. matricariaefolium, Al. Br. Not common.

B. lanceolatum, Angs. Hillside pastures with the preceding two species, and like them very rarely collected.

B. ternatum, Swartz. Pastures and hillsides. Among the numerous varieties described by Prof. D. C. Eaton, the more common are:—
var. intermedium. A large form.

var. obliquum. Medium in size and more common.

var. dissectum. Occasionally with the last. See *Eaton's Ferns of North America*.

B. Virginianum, Swartz. Rich woods; common.

Ophioglossum, L. Adder's-Tongue.

O. vulgatum, L. Bogs and pastures; not uncommon.

ORDER 100. **LYCOPODIACEÆ.** CLUB-MOSS FAMILY.

Lycopodium, L. Club-Moss.

L. lucidulum, Michx. Damp woods; common.

L. inundatum, L. Clayey ground; Grantham Mt., Plymouth, and Hanover, N. H.; Hartland, Vt.; etc.; scarce.

L. annotinum, L. Woods; common.

L. obscurum, L. Seldom distinguished from the variety.

var. *dendroideum*. (L. *dendroideum*, Michx.)—(Tree Club-Moss.) Woods; common.

L. clavatum, L. (Common Club-Moss.) Dry woods; abundant.

L. complanatum, L. (Creeping Ground-Pine.) With the above, common.
var. *chamæcyparissus*, Eaton. Occasional.

ORDER 101. **SELAGINELLACEÆ.** SELAGINELLA FAMILY.

Selaginella, Beauv.

S. rupestris, Spring. Dry rocks; common.

S. apus, Spring. Wet places; about springs; much less common.

Isoetes, L. Quillwort.

I. echinospora, Durieu. The type form is European, but the following varieties are met with:

var. *Braunii*, Engelm. Margin of ponds and streams.

var. *muricata*, Engelm. Mouth of Pompanoosuc river, Norwich, Vt.

var. *Boottii*, Engelm. Goose Pond, Canaan; Conn. river, Hanover.

I. riparia, Engelm. Margin of Conn. river, Hanover.

[I. *lacustris*, L. Assigned to "Lebanon" in the *Geological Survey of New Hampshire*. The species referred to is probably some var. of *I. echinospora*; but if *I. lacustris* is a New England species, it should be found within our limits. Apparently very little is known about it.]

[All the members of this genus are of somewhat uncertain determination. They are found in our springs and ponds, and in the still and shallow margins of our streams, or during a period of drought may be entirely out of the water.]

LIST OF VERTEBRATES.¹

(BY PROF. T. W. D. WORTHEN.)

CLASS I. PISCES. THE FISHES.

ORDER I. NEMATOGNATHI.

FAMILY I. SILURIDÆ. THE CAT-FISHES.

Ameiurus, Rafinesque.

A. nebulosus (Le Sueur). Common Bullhead. Horn Pout. Common.

ORDER II. EVENTOGNATHI.

FAMILY 2. CATOSTOMIDÆ. THE SUCKERS.

Catostomus, Le Sueur. Fine-scaled Suckers.

C. teres (Mitchill). Common Sucker. White Sucker. Abundant.

C. nigricans, Le Sueur. Hog-Sucker. Stone Roller. Common.

FAMILY 3. CYPRINIDÆ. THE MINNOWS.

Notropis, Rafinesque. American Minnows.

N. bifrenatus (Cope). Mud-dace. Brook Minnow. Common.

N. Whipplei (Girard). Silver-Fin. River-Shiner. Not rare.

N. megalops (Rafinesque). Common Shiner. Red-Fin. Dace. Common.

Rhinichthys, Agassiz.

R. atronasus (Mitchill). Black-nosed Dace. "Red-Fin." Common.

Semotilus, Rafinesque.

S. bullaris (Rafinesque). Fall Fish. Chub. Dace. Roach. Abundant.

Notemigonus, Rafinesque.

N. chrysoleucus (Mitchill). Golden Shiner. Shiner. Bream. Abundant.

¹ In this list are included, so far as known, all Animals belonging to the Sub-Kingdom VERTEBRATA, found within about thirty miles of Hanover, N. H. The classification, including citation of authorities, is that given in the *Manual of the Vertebrate Animals of the Northern United States by Pres. David S. Jordan, Fifth edition, rewritten and enlarged*, 1888, where full descriptions may be found.

ORDER III. ISOSPONDYLI. THE SALMON, HERRING, ETC.

FAMILY 4. **SALMONIDÆ.** THE SALMON.**Salvelinus** (Nilsson), Richardson.

- S. namaycush* (Walbaum). Lake Trout. Mackinaw Trout. Salmon Trout. Longe. Rare. *M. V. B. Knox.*
S. fontinalis (Mitchill). Brook Trout. Speckled Trout. Common.
S. aureolus, Bean. Sunapee Lake Trout. In Sunapee Lake. Not rare.

ORDER IV. HAPLOMI. THE PIKE-LIKE FISHES.

FAMILY 5. **ESOCIDÆ.** THE PIKES.**Esox** (Artedi), Linnæus.

- E. americanus*, Gmelin. Grass Pickerel. Common.
E. reticulatus, Le Sueur. Pickerel. Eastern Pickerel. Abundant.
E. lucius, L. Pike. Northern Pickerel. Common.

ORDER V. APODES. THE EELS.

FAMILY 6. **ANGUILLIDÆ.** THE TRUE EELS.**Anguilla**, Thunberg.

- A. anguilla* (L.). Eel. Abundant.

ORDER VI. ACANTHOPTERI. THE SPINY-RAYED FISHES.

FAMILY 7. **CENTRARCHIDÆ.** THE SUN-FISHES.**Lepomis**, Rafinesque. Sun-fishes.

- L. auritus* (Linnaeus). Long-eared Sun-fish. Sun-Perch. Common.
L. gibbosus (L.). Common Sun-fish. Bream. Pumpkin-seed. Abundant.

Micropterus, Lacépède. Black Bass.

- M. dolomieu*, Lacépède. Small-mouthed Black Bass. Abundant.

FAMILY 8. **PERCIDÆ.** THE PERCHES.**Etheostoma**, Rafinesque. Darters.

- E. Olmstedi*, Storer. Darter. Spotted Darter. Common.

Perca (Artedi), Linnaeus.

- P. flavescens* (Mitchill). Yellow-Perch. Perch. Abundant.

Stizostedion, Rafinesque.

- S. vitreum* (Mitchill). Wall-eye. Pike Perch. Plymouth, Vt.

FAMILY 9. **COTTIDÆ.** THE SCULPINS.**Cottus** (Artedi) Linnaeus. Miller's Thumb.

- C. gracilis*, Heckel. Common.

FAMILY 10. **GADIDÆ.** THE COD-FISHES.**Lota**, Cuvier. Burbot. Chub Eel. Ling.

- L. lota* (L.), var. *maculosa*, Le Seur. Eel Pout. Common.

CLASS II. BATRACHIA. THE BATRACHIANS.

ORDER VII. URODELA. THE SALAMANDERS.

FAMILY II. AMBLYSTOMATIDÆ. THE BLUNT-NOSED SALAMANDERS.

Amblystoma, Tschudi.

- A. punctatum* (L.). Spotted Salamander. Common.
A. tigrinum (Green). Not rare.

FAMILY 12. PLETHODONTIDÆ.

Plethodon, Tschudi.

- P. erythronotus* (Green). Common.
P. glutinosus (Green). Not rare.

Spelerves, Rafinesque.

- S. bilineatus* (Green). Not rare.
S. longicauda (Green). Cave Salamander. Rare.
S. ruber (Daudin). Common.

FAMILY 13. PLEURODELIIDÆ. THE NEWTS.

Diemyctylus, Rafinesque.

- D. viridescens*, Rafinesque. Newt. Evet. Eft. Abundant.
 var. *miniatus*, Rafinesque. Red Eft. Common.

ORDER VIII. SALIENTIA. THE TAILLESS BATRACHIANS.

FAMILY 14. BUFONIDÆ. THE TOADS.

Bufo, Laurenti.

- B. lentiginosus*, Shaw. American Toad. Abundant.

FAMILY 15. PELOBATIDÆ. THE BURROWING TOADS.

Scaphiopus, Holbrook.

- S. Holbrooki*, Harlan. Spade-foot. Probable.

FAMILY 16. HYLIDÆ. THE TREE FROGS.

Acris, Dumeril & Bibron.

- A. gryllus*, Le Conte. Cricket Frog. Probable.

Chorophilus, Baird.

C. triseriatus (Wied). Swamp Tree Frog. Probable.

Hyla, Laurenti.

H. versicolor, Le Conte. Common Tree Toad. Common.

H. Pickeringii, Holbrook. Peeping Frog. Common.

FAMILY 17. **RANIDÆ**. THE FROGS.**Rana**, Linnæus.

R. virescens, Kalm. Common Frog. Leopard Frog. Abundant.

R. palustris, Le Conte. Pickerel Frog. Rather common.

R. sylvatica, Le Conte. Wood Frog. Rather common.

R. clamata, Daudin. Green Frog. Common.

R. Catesbeiana, Shaw. Bull Frog. Not rare.

CLASS III. REPTILIA. THE REPTILES.

ORDER IX. OPHIDIA. THE SERPENTS.

FAMILY 18. COLUBRIDÆ. THE COLUBRINE SNAKES.

Storeria, Baird & Girard.

S. occipitomaculata (Storer). Red-Bellied Snake. Common.

Eutainia, Baird & Girard. Garter Snakes.

E. saurita (L.). Riband Snake. Probable.

E. sirtalis (L.). Common Garter Snake. Striped Snake. Abundant.

var. *ordinata* (L.). Common.

var. *dorsalis* (Baird & Girard). Common.

Tropidonotus, Kuhl.

T. sipedon (L.). Water Snake. "Moccasin." Rather rare.

Liopeltis (Fitzinger), Cope.

L. vernalis (De Kay). Grass Snake. Green Snake. Rather common.

Diadophis, Baird & Girard.

D. punctatus (L.). Ring-necked Snake. Not common.

Ophibolus, Baird & Girard.

O. dolius (L.), var. *triangulus* (Boie). Milk Snake. Adder. Spotted Adder. Rather common.

Heterodon, Beauvais.

H. platyrhinus, Latreille. Spreading Adder. Adder. Blowing Viper. Rather common.

FAMILY 19. CROTALIDÆ. THE RATTLESNAKES.

Crotalus, Linnaeus.

C. horridus, L. Common Rattlesnake. Not rare; Springfield, Vt., and vicinity, *C. D. Hull*.

ORDER X. TESTUDINATA. THE TURTLES.

FAMILY 20. CHELYDRIDÆ. THE SNAPPING TURTLES.

Chelydra, Schweigger.

C. serpentina (L.). Common Snapping Turtle. Abundant.

FAMILY 21. **EMYDIDÆ.** THE POND TURTLES.**Chrysemys**, Gray.

C. picta (Hermann). Painted Turtle. Mud Turtle. Common.

Chelopus, Rafinesque.

C. insculptus (Le Conte). Wood Tortoise. Not common.

C. guttatus (Schneider). Speckled Tortoise. Probable.

CLASS IV. AVES. THE BIRDS.

ORDER XI. PYGOPODES. THE DIVING BIRDS.

FAMILY 22. PODICIPIDÆ. THE GREBES.

Podiceps, Latham.

P. cristatus, Latham. Crested Grebe. Rare migrant. *C. H. Storrs.*

Colymbus, Linnaeus.

C. Holbölli (Reinhardt). Red-necked Grebe. Migrant. *C. A. Dowens.*

C. auritus, L. Horned Grebe. Migrant. *C. A. Dowens.*

Podilymbus, Lesson.

P. podiceps (L.). Diedapper. Dab-chick. Dipper Duck. Migrant.

FAMILY 23. URINATORIDÆ. THE LOONS.

Urinator, Cuvier.

U. imber (Gunner). Common Loon. Diver. Northern Diver. Rather common summer resident.

FAMILY 24. ALCIDÆ. THE AUKS.

Alle, Link.

A. alle (L.). Dovekie. Sea Dove. Little Auk. Rare visitor. *F. H. Herrick.*

ORDER XII. LONGIPENNES. THE LONG-WINGED SWIMMERS.

FAMILY 25. LARIDÆ. THE GULLS.

Larus, Linnaeus.

L. argentatus, Brünnich. Herring Gull. Common Gull. Not common summer visitor.

L. Philadelphia, Ord. Bonaparte's Gull. Rare summer visitor.

Sterna, Linnaeus.

S. paradisæa, Brünnich. Arctic Tern. Occasional summer visitor. *M. V. B. Knox.*

S. Antillarum, Lesson. Least Tern. Rare summer visitor.

FAMILY 26. PHALACROCORACIDÆ. THE CORMORANTS.

Phalacrocorax, Brisson.

P. carbo (L.). Common Cormorant. Very rare visitor. *C. A. Dowens.*

ORDER XIII. ANSERES. THE DUCKS AND GEESE.

FAMILY 27. ANATIDÆ. THE DUCKS AND GEESE.

Merganser, Brisson.

M. Americanus (Cassin). Merganser. Goosander. Not rare migrant.

Lophodytes, Reichenbach.

L. cucullatus (L.). Hooded Merganser. Sheldrake. Occasional visitor.
M. V. B. Knox.

Anas, Linnæus.

- A. *boschas*, L. Mallard Duck. Tame Duck. "Original of the common domesticated duck." Rather rare migrant.
- A. *obscura*, Gmelin. Black Duck. Common migrant. Occasional summer resident.
- A. *discors*, L. Blue-winged Teal. Migrant.
- A. *Carolinensis*, Gmelin. Green-winged Teal. Migrant.

Aix, Boie.

A. sponsa (L.). Wood Duck. Summer Duck. Not rare summer resident.

Erismatura, Bonaparte.

E. rubida (Wilson). Ruddy Duck. Common migrant.

Branta, Scopoli.

B. Canadensis (L.). Wild Goose. Canada Goose. Rather common migrant.

ORDER XIV. HERODIONES. THE HERONS AND STORKS.

FAMILY 28. ARDEIDÆ. THE HERONS.

Botaurus, Stephens.

- B. lentiginosus* (Montagu). Bittern. Indian Hen. Common summer resident.
- B. exilis* (Gmelin). Least Bittern. Not rare summer resident. *C. H. Storrs.*

Ardea, Linnæus.

- A. herodias*, L. Great Blue Heron. Rather common summer resident.
- A. virescens* (L.). Green Heron. Rather common summer resident.

Nycticorax, Stephens.

- N. nycticorax* (L.). Night-heron. Qua Bird. Squawk. Rather rare summer resident.

ORDER XV. PALUDICOLÆ. THE CRANES AND RAILS.

FAMILY 29. RALLIDÆ. THE RAILS.

Rallus, Linnæus.

R. Virginianus, L. Virginia Rail. Probable summer resident.

Porzana, Vieillot.

P. Carolina (L.). Carolina Rail. Sora. Occasional summer resident.
R. Chellis.

Fulica, Linnæus.

F. Americana (Gmelin). Coot. Mud Hen. Rather rare summer resident.

ORDER XVI. LIMICOLÆ. THE SHORE-BIRDS.

FAMILY 30. SCOLOPACIDÆ. THE SNIPE.

Philohela, Gray.

P. minor (Gmelin). American Woodcock. Common migrant; not rare summer resident.

Gallinago, Leach.

G. delicata (Ord.). Wilson's Snipe. Not rare summer resident. *C. H. Storrs*; *R. Chellis.*

Tringa, Linnæus.

T. maritima, Brünnich. Purple Sandpiper. Not common summer visitor.
T. minutilla, Vieillot. Least Sandpiper. Peep. Rare summer visitor. *C. H. Storrs.*

Totanus, Bechstein.

T. melanoleucus (Gmelin). Greater Tell-tale. Yellow Shanks. Occasional summer visitor. *C. H. Storrs*; *R. Chellis.*

T. flavipes (Gmelin). Lesser Tattler. Yellow Legs. Not common summer visitor.

Bartramia, Lesson.

B. longicauda (Bechstein). Upland Sandpiper. Rather common summer resident.

Actitis, Boie.

A. macularia (L.). Tip-up. Teeter-Tail. Spotted Sandpiper. Common summer resident.

FAMILY 31. CHARADRIIDÆ. THE PLOVERS.

Ægialitis, Boie.

A. vocifera (L.). Kildeer. Rare summer visitor. *B. T. Blanpied.*
A. meloda (Ord.). Piping Plover. Occasional summer resident. *C. H. Storrs.*

Arenaria, Brisson.

A. interpres (L.). Turnstone. Very rare summer visitor. *C. A. Downs.*

ORDER XVII. GALLINÆ. THE GALLINACEOUS BIRDS.

FAMILY 32. TETRAONIDÆ. THE GROUSE.

Colinus, Lesson.

C. Virginianus (L.). Bob-White. Quail. Rare summer visitor or resident. *T. H. Chubb*; *B. T. Blanpied.*

Dendragapus, Elliott.

D. Canadensis (L.). Spruce Partridge. Canada Grouse. Very rare resident.

Bonasa, Stephens.

B. umbellus (L.). Ruffed Grouse. Partridge. Common resident.

ORDER XVIII. COLUMBÆ. THE DOVES.

FAMILY 33. **COLUMBIDÆ**. THE PIGEONS.**Ectopistes**, Swainson.

E. migratorius (L.). Wild Pigeon. Passenger Pigeon. Rather rare summer resident.

Zenaidura, Bonaparte.

Z. macroura (L.). Mourning Dove. Turtle Dove. Carolina Dove. Very rare summer resident.

ORDER XIX. RAPTORES. THE BIRDS OF PREY.

FAMILY 34. **FALCONIDÆ**. THE FALCONS.**Circus**, Lacépède.

C. Hudsonius (L.). Marsh Harrier. Common summer resident.

Accipiter, Brisson.

A. velox (Wilson). Sharp-Shinned Hawk. "Pigeon Hawk." Rather common summer resident.

A. Cooperi, Bonaparte. Chicken Hawk. Common summer resident.

A. atricapillus (Wilson). Goshawk. Rather rare winter visitor and rare resident.

Buteo, Cuvier.

B. borealis (Gmelin). Hen Hawk. Red-Tailed Hawk. Common resident.

B. latissimus (Wilson). Broad-Winged Hawk. Not rare summer resident.

B. lineatus (Gmelin). Chicken Hawk. Red-Shouldered Hawk. Rather common resident.

Archibuteo, Brehm.

A. lagopus (Brünnich). Rough-Legged Hawk. Black Hawk. Probable resident.

Aquila, Brisson.

A. chrysaëtos (L.). Golden Eagle. Probable summer resident. *Knox*.

Haliaëtus, Savigny.

H. leucocephalus (L.). Bald Eagle. Rather rare summer resident.

Falco, Linnaeus. Falcons.

F. rusticulus (L.). Gray Gyrfalcon. Jerfalcon. Probable winter visitor.

F. peregrinus, Tunstall. Peregrine Falcon. Duck Hawk. Not common resident.

F. columbarius, L. Pigeon Hawk. American Merlin. Rather common summer resident and rare resident.

F. sparverius, L. Sparrow Hawk. Rusty Crowned Falcon. Rather common summer resident.

Pandion, Savigny.

P. haliaetus (L.). Osprey. Fish Hawk. Rather common summer resident.

FAMILY 35. **BUBONIDÆ. THE OWLS.**

Asio, Brisson.

A. Wilsonianus (Lesson). Long-eared Owl. Rare resident. *T. Bakes.*

A. accipitrinus (Pallas). Short-eared Owl. Very rare resident.

Syrnium, Savigny.

S. nebulosum (Forster). Barred Owl. Rather common resident.

Scotiaptex, Swainson.

S. cinereum (Gmelin). Great Gray Owl. Spectral Owl. Probable winter visitor.

Nyctala, Brehm.

N. Tengmalmii (Gmelin). Sparrow Owl. Probable winter visitor.

N. Acadica (Gmelin). Saw-whet Owl. Acadian Owl. Rather rare resident.

Megascops, Kaup.

M. asio (L.). Screech Owl. Red Owl. Rather common resident.

Bubo, Duméril.

B. Virginianus (Gmelin). Great Horned Owl. Not rare resident.

Nyctea, Stephens.

N. nyctea (L.). Snowy Owl. Very rare winter visitor.

Surnia, Duméril.

S. ulula (L.). Hawk Owl. Day Owl. Rare winter visitor or resident.

ORDER XX. COCCYGES. THE CUCKOO-LIKE BIRDS.

FAMILY 36. **CUCULIDÆ. THE CUCKOOS.**

Coccyzus, Vieillot.

C. Americanus (L.). Yellow-billed Cuckoo. "Rain Crow." Rather common summer resident,

C. erythrophthalmus (Wilson). Black-billed Cuckoo. Common summer resident.

FAMILY 37. **ALCEDINIDÆ. THE KINGFISHERS.**

Ceryle, Boie.

C. alcyon (L.). Belted Kingfisher. Common summer resident.

ORDER XXI. PICI. THE WOODPECKERS AND WRYNECKS.

FAMILY 38. **PICIDÆ.** THE WOODPECKERS.**Dryobates**, Boie.

- D. villosus* (L.). Hairy Woodpecker. Big Sap-Sucker. Common winter visitor; not rare resident.
D. pubescens (L.). Downy Woodpecker. Little Sap-Sucker. Common resident.

Picoides, Lacépède.

- P. arcticus* (Swainson). Black-backed Woodpecker. Not common winter resident.
P. Americanus, Brehm. Striped Woodpecker. Not rare winter visitor.

Sphyrapicus, Baird.

- S. varius* (L.). Yellow-bellied Woodpecker. Common summer resident.

Ceophleæus, Cabanis.

- C. pileatus* (L.). Logcock. Big Black Woodpecker. Pileated Wood-pecker. Not rare resident.

Melanerpes, Swainson.

- M. erythrocephalus* (L.). Red-headed Woodpecker. Rather rare summer resident.
M. Carolinus (L.). Red-bellied Woodpecker. Possible summer resident.

Colaptes, Swainson.

- C. auratus* (L.). Yellow-Hammer. Flicker. Wood-wall. Golden-winged Woodpecker. High-Holer. Common summer resident.

ORDER XXII. MACROCHIRES. THE SWIFTS AND HUMMING BIRDS.

FAMILY 39. **CAPRIMULGIDÆ.** THE GOATSUCKERS.**Antrostomus**, Gould.

- A. vociferus* (Wilson). Whippoorwill. Night Jar. Common summer resident.

Chordeiles, Swainson.

- C. Virginianus* (Gemlin). Night Hawk. Bull Bat. Very common summer resident.

FAMILY 40. **MICROPODIDÆ.** THE SWIFTS.**Chætura**, Stephens.

- C. pelágica* (L.). Chimney Swift. Chimney Swallow. Abundant summer resident.

FAMILY 41. **TROCHILIDÆ.** THE HUMMING BIRDS.**Trochilus**, Linnæus.

- T. colubris*, L. Ruby-throated Humming Bird. Common summer resident.

ORDER XXIII. PASSERES. THE PASSERINE BIRDS.

FAMILY 42. TYRANNIDÆ. THE FLYCATCHERS.

Tyrannus, Cuvier.

T. tyrannus (L.). King Bird. Bee Martin. Common summer resident.

Myiarchus, Cabanis.

M. crinitus (L.). Great Crested Flycatcher. Not rare summer resident.

Sayornis, Bonaparte.

S. Phœbe (Latham). Pewee. Phœbe. Abundant summer resident.

Contopus, Cabanis.

C. borealis (Swainson). Olive-sided Flycatcher. Rather rare summer resident.

C. virens (L.). Wood Pewee. Common summer resident.

Empidonax, Cabanis.

E. flaviventris, Baird. Yellow-bellied Flycatcher. Not common summer resident.

E. Acadianus (Gmelin). Small Green-crested Flycatcher. Not rare summer resident.

E. pusillus (Swainson), var. *Trailli*, Audubon. Traill's Flycatcher. Not common summer resident.

E. minimus, Baird. Least Flycatcher. Very common summer resident.

FAMILY 43. ALAUDIDÆ. THE LARKS.

Otocoris, Bonaparte.

O. alpestris (Forster). Shore Lark. Horned Lark. Not rare migrant.

G. D. Hull.

FAMILY 44. CORVIDÆ. THE CROWS AND JAYS.

Cyanocitta, Strickland.

C. cristata (L.). Blue Jay. Common resident.

Perisoreus, Bonaparte.

P. Canadensis (L.). Canada Jay. Gray Jay. Whiskey Jack. Occasional visitor; possible resident. *Storrs.*

Corvus, Linnæus.

C. Americanus, Audubon. Crow. Very common resident.

FAMILY 45. ICTERIDÆ. THE AMERICAN "ORIOLES" AND "BLACK-BIRDS."

Dolichonyx, Swainson.

D. oryzivorus (L.). Bobolink. Reed Bird. Rice Bird. Very common summer resident.

Molothrus, Swainson.

M. ater (Boddaert). Cow Blackbird. Cowbird. Common summer resident.

Agelaius, Vieillot.

A. *phoeniceus* (L.). Red-winged Blackbird. Swamp Blackbird. Common summer resident.

Sturnella, Vieillot.

S. *magna* (L.). Meadowlark. Not rare summer resident.

Icterus, Brisson.

I. *galbula* (L.). Baltimore Oriole. Golden Robin. Fire Bird. Common summer resident.

Scolecophagus, Swainson.

S. *Carolinus* (Müller). Rusty Grackle. Rusty Blackbird. Not rare migrant.

Quiscalus, Vieillot.

Q. *quiscula* (L.). Crow Blackbird. Purple Grackle. Rather common summer resident.

FAMILY 46. **FRINGILLIDÆ. THE FINCHES.****Pinicola**, Vieillot.

P. *enucleator* (L.). Pine Grosbeak. Rather common winter resident.

Carpodacus, Kaup.

C. *purpureus* (Gmelin). Purple Finch. Very common summer resident.

Loxia, Linnæus.

L. *leucoptera*, Gmelin. White Winged Crossbill. Rather rare winter visitor.

L. *curvirostra* (L.). Red Crossbill. Not rare winter resident; rare resident.

Acanthis, Bechstein.

A. *linarius* (L.). Red Poll Linnet. Rather common winter resident.

Spinus, Boie.

S. *tristis* (L.). Yellow Bird. Thistle Bird. American Goldfinch. Abundant resident.

S. *pinus* (Wilson). Pine Siskin. Pine Finch. Rather common winter visitor; not rare resident.

Plectrophenax, Stejneger.

P. *nivalis* (L.). Snow Bunting. Very common winter resident.

Calcarius, Bechstein.

C. *Lapponicus* (L.). Lapland Longspur. Rare winter visitor.

Poocætes, Baird.

P. *gramineus* (Gmelin). Bay-winged Bunting. Grass Sparrow. Ground Bird. Abundant summer resident.

Passer, Brisson.

P. *domesticus* (L.). English Sparrow. Abundant resident.

Ammodramus, Swainson.

A. *Sandwichensis* (Gmelin), var. Savanna. Savanna Sparrow. Not common summer resident.

A. savannarum (Gmelin). Grasshopper Sparrow. Probable summer resident.

Zonotrichia, Swainson.

Z. leucophrys (Forster). White-crowned Sparrow. Common migrant.

Z. albicollis (Gmelin). White-throated Sparrow. Peabody Bird. Rather common resident.

Spizella, Bonaparte.

S. monticola (Gmelin). Tree Sparrow. Rather rare winter visitor.

S. socialis (Wilson). Chippy. Chipping Sparrow. Abundant summer resident.

S. pusilla (Wilson). Field Sparrow. Common summer resident.

Junco, Wagler.

J. hyemalis (L.). Snow Bird. Common migrant; not rare resident.

Melospiza, Baird.

M. fasciata (Gmelin). Song Sparrow. Very common summer resident.

M. Georgiana (Latham). Swamp Sparrow. Rather common summer resident.

Passerella, Swainson.

P. iliaca (Merrem). Fox Sparrow. Common migrant.

Pipilo, Vieillot.

P. erythrophthalmus (L.). Chewink. Marsh Robin. Towhee. Not rare summer resident.

Habia, Reichenbach.

H. Ludoviciana (L.). Rose-breasted Grosbeak. Rather common summer resident.

Passerina, Vieillot.

P. cyanea (L.). Indigo Bird. Very common summer resident.

FAMILY 47. **TANAGRIDÆ**. THE TANAGERS.

Piranga, Vieillot.

P. erythromelas, Vieillot. Scarlet Tanager. Rather common summer resident.

FAMILY 48. **HIRUNDINIDÆ**. THE SWALLOWS.

Progne, Boie.

P. subis (L.). Purple Martin. Rather common summer resident.

Petrochelidon, Cabanis.

P. lunifrons (Say). Cliff Swallow. Abundant summer resident.

Chelidon, Forster.

C. erythrogaster (Boddaert). Barn Swallow. Abundant summer resident.

Tachycineta, Cabanis.

T. bicolor (Vieillot). White-bellied Swallow. Very common summer resident.

Clivicola, Forster.

C. riparia, L. Bank Swallow. Sand Martin. Very common summer resident.

FAMILY 49. **AMPELIDÆ**. THE CHATTERERS.**Ampelis**, Linnæus.

A. garrulus, L. Bohemian Wax Wing. Northern Wax Wing. Probable winter visitor.

A. cedrorum (Vieillot). Cedar Bird. Cherry Bird. Very common summer resident.

FAMILY 50. **LANIIDÆ**. THE SHRIKES.**Lanius**, Linnæus.

L. borealis, Vieillot. Great Northern Shrike. Butcher-bird. Rather common winter resident, and rather rare resident.

L. Ludovicianus, L. Logger-head Shrike. Rather rare resident.

FAMILY 51. **VIREONIDÆ**. THE VIREOS.**Vireo**, Vieillot.

V. olivaceus (L.). Red-eyed Vireo. Greenlet. Very common summer resident.

V. gilvus (Vieillot). Warbling Vireo. Rather common summer resident.

V. flavifrons, Vieillot. Yellow-throated Vireo. Not rare summer resident.

V. solitarius (Wilson). Blue-headed Vireo. Rather rare summer resident.

V. Noveboracensis (Gmelin). White-eyed Vireo. Rather common summer resident.

FAMILY 52. **MNIOTILTIDÆ**. THE NEW WORLD WARBLERS.**Mniotilla**, Vieillot.

M. varia (L.). Black and White Creeper. Rather common summer resident.

Helminthophila, Ridgway.

H. chrysoptera (L.) Golden-winged Warbler. Probable migrant and resident.

H. pinus (L.). Blue-winged Yellow Warbler. Possible summer resident.

H. peregrina (Wilson). Tennessee Warbler. Rare migrant, and possible resident.

H. celata (Say). Orange-crowned Warbler. Rare summer resident. *F. H. Herrick*.

H. ruficapilla (Wilson). Nashville Warbler. Not common summer resident.

Compsothlypis, Cabanis.

C. Americana (L.). Blue Yellow-backed Warbler. Not common summer resident.

Dendroica, Gray.

D. aestiva (Gmelin). Summer Warbler. Golden Warbler. Rather common summer resident.

D. cærulescens (L.). Black-throated Blue Warbler. Rather rare summer resident. *C. H. Storrs.*

D. coronata (L.) Yellow-rumped Warbler. Common migrant and probable resident.

D. maculosa (Gmelin). Black and Yellow Warbler. Not rare summer resident.

D. Pennsylvanica (L.). Chestnut-sided Warbler. Rather common summer resident.

D. castanea (Wilson). Bay-breasted Warbler. Autumn Warbler. Rare summer resident.

D. striata (Forster). Black-poll Warbler. Common migrant.

D. Blackburniæ (Gmelin). Orange-throated Warbler. Blackburnian Warbler. Rare summer resident.

D. virens (Gmelin). Black-throated Green Warbler. Rather common summer resident.

D. vigorsi (Audubon). Pine-creeping Warbler. Common migrant, and rather rare summer resident.

D. discolor (Vieillot). Prairie Warbler. Probable summer resident.

D. palmarum (Gmelin). Red-poll Warbler. Rather common migrant.

Seiurus, Swainson.

S. aurocapillus (L.). Oven-bird. Golden-crowned "Thrush." Rather common summer resident.

S. Noveboracensis (Gmelin). Water Wagtail. Water Thrush. Probable summer resident.

Geothlypis, Cabanis.

G. Philadelphia (Wilson). Mourning Warbler. Rare summer resident.

G. trichas (L.) Maryland Yellow Throat. Common summer resident.

Sylvania, Nuttall.

S. pusilla (Wilson). Green Black-capped Warbler. Rather common migrant.

S. Canadensis (L.). Canada Warbler. Not common summer resident. *C. H. Storrs.*

Setophaga, Swainson.

S. ruticilla (L.). American Redstart. Rather common summer resident.

FAMILY 53. **MOTACILLIDÆ. THE WAGTAILS.****Anthus**, Bechstein.

A. Pennsylvanicus (Latham). Brown Lark. Tit-lark. Rather common migrant.

FAMILY 54. **TROGLODYTIDÆ.** THE WRENS AND MOCKING-BIRDS.

Mimus, Boie.

M. polyglottos (L.). Mocking-Bird. Very rare visitor. *C. A. Dowens.*

Galeoscoptes, Cabanis.

G. Carolinensis (L.). Cat-Bird. Very common summer resident.

Harporrynchus, Cabanis.

H. rufus (L.). Brown Thrush. Brown Thrasher. Rather common summer resident.

Troglodytes, Vieillot.

T. aedon, Vieillot. House Wren. Common summer resident.

T. hiemalis, Vieillot. Winter Wren. Not rare migrant.

Cistothorus, Cabanis.

C. stellaris (Lichtenstein). Short-billed Marsh Wren. Probable summer resident.

C. palustris (Wilson). Long-billed Marsh Wren. Possible summer resident.

FAMILY 55. **CERTHIIDÆ.** THE CREEPERS.

Certhia, Linnæus.

C. familiaris, L. Brown Creeper. Rather rare summer resident.

FAMILY 56. **PARIDÆ.** THE NUTHATCHES AND TITMICE.

Sitta, Linnæus.

S. Carolinensis, Latham. White-bellied Nuthatch. "Sap-sucker." Common resident.

S. Canadensis, L. Red-bellied Nuthatch. Common winter resident, and rather rare resident.

Parus, Linnæus.

P. atricapillus, L. Black-capped Titmouse. Chickadee. Very common resident.

FAMILY 57. **SYLVIIDÆ.** THE OLD WORLD WARBLERS.

Regulus, Cuvier.

R. satrapa, Lichtenstein. Golden-crowned Kinglet. Common migrant, and not rare winter resident.

R. calendula (L.). Ruby-crowned Kinglet. Common migrant.

FAMILY 58. **TURRIDÆ.** THE THRUSHES.

Turdus, Linnæus.

T. mustelinus, Gmelin. Wood Thrush. Rare summer resident.

T. fuscescens, Stephens. Tawny Thrush. Wilson's Thrush. Not common summer resident.

T. ustulatus, Nuttall. Olive-backed Thrush. Rare summer resident.

T. Aonalaschkæ, Gmelin, var. Pallasi, Cabanis. Hermit Thrush. Common summer resident.

Merula, Leach.

M. migratoria (L.). Robin. American Red Breast. Abundant summer resident, and very rare resident.

Sialia, Swainson.

S. sialis (L.). Common Blue Bird. Very common summer resident.

CLASS V. MAMMALIA. THE MAMMALS.

ORDER XXIV. GLIRES. THE RODENTS OR GNAWERS.

FAMILY 59. **LEPORIDÆ.** THE HARES.**Lepus**, Linnæus.

L. Americanus, Erxleben. White Rabbit. Northern Hare. Common.

FAMILY 60. **HYSTRICIDÆ.** THE PORCUPINES.**Erethizon**, Frédéric Cuvier.

E. dorsatus (L.). Canada Porcupine. Not rare. This is wrongly called "Hedgehog", which is European.

FAMILY 61. **ZAPOLIDÆ.** THE JUMPING MICE.**Zapus**, Coues.

Z. Hudsonius (Zimmerman). Jumping Mouse. Rare.

FAMILY 62. **MURIDÆ.** THE MICE.**Fiber**, Cuvier.

F. zibethicus (L.). Muskrat. Common.

Arvicola, Lacépède. Field Mice.

A. Pennsylvanicus, Ord. Meadow Mouse. Abundant.

Evotomys, Coues.

E. rutilus (Pallas). Long-eared Mouse. Rather rare.

Calomys, Waterhouse.

C. Americanus (Kerr). Common White-footed Mouse. Dormouse. Common.

Mus, Linnæus. Old World Rats and Mice.

M. decumanus, Pallas. Brown or Norway Rat. Abundant.

M. rattus, L. Black Rat. Common.

M. musculus, L. Common House Mouse. Too abundant.

FAMILY 63. **CASTORIDÆ.** THE BEAVERS.**Castor**, Linnæus.C. fiber, L. Beaver. Very rare. *C. A. Downs.*

FAMILY 64. SCIURIDÆ. THE SQUIRRELS.

Arctomys, Schreber.

A. monax (L.). Woodchuck. Ground Hog. Very common.

Tamias, Illiger.

T. striatus (L.). Chipmunk. Ground Squirrel. Abundant.

Sciurus, Linnæus.

S. Hudsonicus, Erxleben. Red Squirrel. Chickaree. Very common.

S. Carolensis, Gmelin. Includes,—Gray Squirrel, common; Black Squirrel, very rare.

Sciuropterus, Frédéric Cuvier.

S. volans (L.). Common Flying Squirrel. Rather common.

ORDER XXV. INSECTIVORA. THE INSECT-EATERS.

FAMILY 65. SORICIDÆ. THE SHREWS.

Sorex, Linnæus.

S. Forsteri (Richardson). Probable.

S. platyrhinus (De Kay). Common Shrew. Shrew Mouse. Not rare.

Blarina, Gray.

B. brevicauda (Say). Mole-Shrew. Probable.

B. angusticeps, Baird. Not rare.

FAMILY 66. TALPIDÆ. THE MOLES.

Scalops, Cuvier,

S. aquaticus (L.). Common Mole. Common.

Scapanus, Pomel.

S. Americanus (Bartram). Hairy-tailed Mole. Probable.

Condylura, Illiger.

C. cristata (L.). Star-nosed Mole. Common.

ORDER XXVI. CHEIROPTERA. THE BATS.

FAMILY 67. VESPERTILIONIDÆ. THE COMMON BATS.

Vespertilio, Linnæus.

V. subulatus, Say. Little Brown Bat. Very common.

Vesperugo, Keyserling & Blasius.

V. noctivagans (Le Conte). Silver Black Bat. Not common. *R. Chellis*.

V. Georgianus (F. Cuvier). Rare.

V. serotinus (Schreber), var. *fuscus*, Beauvais. Long-eared Bat. Rare.

Atalapha, Rafinesque.

A. Noveboracensis (Erxleben). Red Bat. Common.

A. cinerea (Beauvais). Hoary Bat. Possible.

ORDER XXVII. UNGULATA. THE HOOFED MAMMALS.

FAMILY 68. CERVIDÆ. THE DEER.

Cariacus, Gray.

C. Virginianus (Boddaert). Virginia Deer. Red Deer. Rare. Has been reintroduced into Vermont, and may be seen occasionally throughout the district.

ORDER XXVIII. FERÆ. THE FLESH-EATERS OR CARNIVORA.

FAMILY 69. PROCYONIDÆ. THE RACCOONS.

Procyon, Storr.

P. lotor (L.). Common Raccoon. Coon. Rather common.

FAMILY 70. URSIDÆ. THE BEARS.

Ursus, Linnæus.

U. Americanus, Pallas. Brown, Black, or Cinnamon Bear. Rare.

FAMILY 71. MUSTELIDÆ. THE WEASELS.

Mephitis, Cuvier.

M. mephitica (Shaw). Common Skunk. Too common.

Gulo, Storr.

G. gulo (L.). Wolverine. Probable.

Mustela, Linnæus.

M. Americana, Turton. Sable. Pine Marten. Not common.

M. Pennanti (Erxleben). Black Cat. Rare.

Putorius, Cuvier.

P. vison (Schreber). Mink. Rather common.

P. erminea (L.). Weasel. Ermine. Stoat. Rather common.

P. nivalis (L.). Least Weasel. Rare.

FAMILY 72. CANIDÆ. THE DOGS.

Vulpes, Brisson.

V. vulpes (L.). Red Fox. Rather common.

var. *decussatus*, Desm. The Cross Fox. Very rare.

var. *argentatus*, Shaw. Black or Silver Fox. Very rare.

FAMILY 73. FELIDÆ. THE CATS.

Lynx, Rafinesque.

L. Canadensis (Desmarest). Canada Lynx. Rare.

L. rufus (Guldenstädt). American Wild Cat. Rare.

Felis, Linnæus. Cats.

F. concolor, L. American Panther. Cougar. Puma. Very rare.

ORDER XXIX. PRIMATES. THE ANTHROPOID MAMMALS.

FAMILY 74. HOMINIDÆ. THE MEN.

Homo, Linnæus.

- H. sapiens, L. Man.
 var. Americanus, L. "American Indian."
 var. Europæus, L. Caucasian race.
 var. Asiaticus, L. Mongolian race.
 var. Afer, L. Negro race.

SUMMARY OF SPECIES.

Fishes,	24
Batrachians,	20
Reptiles,	15
Birds,	191
Mammals,	50
Total,	<hr/> 300

NOTE.—The above List, it is hoped, will furnish a good working basis for a thorough exploration of the region designated, and ultimately for a complete Catalogue of the Vertebrates occurring within our limits. The coöperation of all naturalists and sportsmen is invited, and they are requested to report all additions, corrections, and information of any kind, to either PROF. T. W. D. WORTHEN or PROF. H. G. JESUP, Hanover, N. H.

INDEX TO PLANTS.

[SYNONYMS IN PARENTHESES; CULTIVATED PLANTS IN ITALICS.]

Abies	41	Apple of Peru	30	Beard-tongue	30
Abutilon	7	(Aquitoliaceæ)	8	Bedstraw	20
Acalypha	36	Aquilegia	2	Bee-balm	32
Acer	9	Arabis	4	Beech-drops	31
Achillea	23	Araceæ	47	Beech	39
Aconitum	2	Arachis	12	Blue	38
Acorus	47	Aralia	17	Beet,	34
Actaea	2	Araliaceæ	17	Beggar's-lice	28
Adder's-mouth	42	Arbor-vitæ	41	Bellflower	25
Adder's-tongue	45, 58	Arbutus, Trailing	26	Bellwort	45
Adder's-tongue Family	58	Arctothibium	36	Berberidaceæ	2
Adiantum	56	Aretium	23	Berberis	2
Adlumia	8	Arctostaphylos	25	Bergamot	32
Æsculus	9	Arenaria	6	Beta	34
Agrimonias	13	Arethusa	43	Betula	38
Agropyrum	55	Arisema	47	(Betulaceæ)	38
Agrostis	53	Aristida	52	Bidens	22
(Aira)	54	Aristolochia	36	Bindweed	29, 35
Alder	38	Aristolochiaceæ	36	Birch Family	38
Black	8	Arrhenatherum	53	Birthwort	36, 45
Alisma	47	Arrow-head	47	Birthwort Family	36
Alismaceæ	47	Artemisia	23	Bishop's-cap	14
Allium	44	Arum, Arrow	47	Bitter-put	38
Alnus	38	Water	47	Bittersweet	29
Alopecurus	53	Arum Family	47	Climbing	8
Alysseum	5	Asarum	36	Blackberry	12
Amarantaceæ	34	Asclepiadaceæ	27	Bladder-nut	10
Amaranth Family	34	Asclepias	27	Bladderwort Family	31
Amarantus	34	Ash	27	Blephilia	33
Ambrosia	22	Mountain	13	Blite, Strawberry	34
Amelanchier	14	Prickly	8	(Blitum)	34
Ampelopsis	9	Asparagus	45	Blood-root	3
Amphicarpa	11	Aspen	39	Blueberry	25
Anacardiaceæ	10	Asphodel, False	46	Blue-curls	32
Anagallis	27	Aspidium	57	Blue-flag	44
Anaphalis	22	Asplenium	56	Bluet	19
Andromeda	26	Asprella	55	Boehmeria	37
Andropogon	52	Aster	21	Boneset	20
Anemone	1, 2	Astragalus	11	Borage Family	28
Anemonella	1	Avena	54	Borraginaceæ	28
Angelica	16	Avens	12	Botrychium	58
Antennaria	22	(Azalea)	26	Boxelder	9
Anthemis	23			Brachelytrum	53
Anthoxanthum	52	Balm of Gilead	40	Brake	56
Anychia	34	Balsam	8	Cliff	56
Aphyllon	31	Baneberry	2	Brasenia	3
Apios	11	Barbarea	4	Brassica	4, 5
Apium	17	Barberry	2	Bromus	55
Aplectrum	42	Barberry Family	2	Broom-rape Family	31
Apocynaceæ	27	Barley	55	Brunella	33
Apocynum	27	Basil	32, 33	Buckbean	28
Appéle	14	Basswood	7	Buckeye	9
Balsam	16	Bean	12	Buckthorn	9
Crab	14	Bearberry	25	Buckthorn Family	9

Buckwheat	35	Chess	55	Cucumber	16
Buckwheat Family	35	Chestnut	39	Bur	16
Buda	6	Chickweed	6	Cucumber Family	16
Bulrush	49	Forked	34	Cucumber-root, Indian	45
Bunch-berry	17	Chicory	24	Cucumis	16
Burdock	23	Chimaphila	26	Cucurbita	16
Burnet	13	Chiogenes	25	Cucurbitaceæ	16
Burning-bush	8	Chiouanthus	27	Cudweed	22
Bur-reed	47	Chives	44	Cupuliferæ	38
Buttercup	2	Choke-berry	13	Currant	14
Butternut	37	Chrysanthemum	23	Cusentia	29
Button-bush	20	Chrysopogon	52	Cyclonia	14
Buttonwood	37	Chrysosplenium	14	Cynoglossum	28
		Cichorium	24	Cyperaceæ	48
<i>Cabbage</i>	5	Cicta	17	Cyperus	48
Cabbage, Skunk	47	Cimicifuga	2	Cypress-vine	29
Calamagrostis	53	Cinna	53	Cypripedium	44
Calamintha	32	Cinque-foil	13	Cystopteris	57
Calla	47	Ciræa	16	Dactylis	54
Callitrichæ	15	(Cirsium)	24	Daisy, Ox-eye	28
Calopogon	43	Cistaceæ	5	Dalibarda	12
Caltha	2	Citrulus	16	Dandelion	24
Calypso	42	Cladium	49	Danthonia	54
(Calystegia)	29	Claytonia	7	Daphne	36
Camelina	4	Clearweed	37	Datura	30
Campanulaceæ	25	Clematis	1, 2	Daucus	16
Campanula Family	25	Clintonia	45	Delphinium	2
Camptosorus	57	Clover	10	Dentaria	4
Cancer-root	31	Bush	11	Deschampsia	54
Cloudytuft	5	Sweet	10	Desmodium	11
Cannabis	37	Club-moss	58	Dianthus	6
Caprifoliaceæ	19	Club-moss Family	58	Dicentra	3
Capsella	4	Cnicus	24	Dicksonia	57
Caragana	12	Cocklebur	22	Diervilla	19
Caraway	17	Cockscomb	34	Direa	36
Cardamine	4	Cohosh, Black	2	Dock	35
Cardinal-flower	25	Blue	2	Dodder	29
Carex	49-51	Colchicum	46	Dogbane Family	27
Carpet-weed	16	Coltsfoot	23	Dogwood Family	17
Carpinus	38	Sweet	23	Dragon-head	33
Carrot	16	Columbine	2	Drosera	15
Carum	17	Comandra	36	Droseraceæ	15
Carya	38	Comfrey	28, 29	Duckweed Family	47
Caryophyllaceæ	6, 34	Compositæ	20	Dulichium	49
Cashew Family	10	Composite Family	20	Dutchman's-breeches	3
Cassandra	26	Cone-flower	22		
Cassia	11	Coniferae	41	Eatonia	54
Castanea	39	Conioselinum	16	Echinocystis	16
Castor-oil plant	37	Conium	17	Echinospermum	28
Catchfly	6	Conopholis	31	Echinum	29
Catnip	33	Convallaria	44	Elder	19
Cat-tail Family	47	Convolvulaceæ	29	Elecampane	22
Caulophyllum	2	Convolvulus Family	29	Eleocharis	49
Ceanothus	9	Coptis	2	Elm	37
Cedar, White	41	Corallorrhiza	42	Elodea	42
Red	41	Coral-root	42	Elodes	7
Celandine	3	Cornaceæ	17	Elymus	55
Celastraceæ	8	Corn-cockle	6	Empetraceæ	40
Celastrus	8	Cornel	17, 18	Empetrum	40
Celery	17	Cornus	17	Epigaea	26
Celosia	34	Coronilla	12	Epilobium	15
Celtis	37	Corydalis	3	Epiphegus	31
Cenchrus	52	Corylus	38	Equisetaceæ	56
Cephalanthus	20	Cottonwood	40	Equisetum	56
Cerastium	6	Cowslip	25	Eragrostis	54
Ceratophyllaceæ	40	Cranberry	19	Erechitites	23
Ceratophyllum	40	High	19	Ericacæ	25
Chamomile	23	Crassulaceæ	15	Erigeron	21
Checkerberry	26	Crataegus	13	Eriocaulæ	48
Chelidonium	3	Cress	4	Eriocaulon	48
Cyclone	30	Crocus	44	Eriophorum	49
Chenopodiaceæ	34	Crowberry Family	40	Erodium	8
Chenopodium	34	Crowfoot Family	1	Erysimum	4
Cherry	12	Cruciferae	4	Erythronium	45
Ground	29	Cryptotaenia	17	Euonymus	8

Eupatorium	20	Gomphrena	34	Helianthus	22
Euphorbia	36	Good-King-Heury	34	Heliopsis	22
Euphorbiaceæ	36	Goodyera	43	Hellebore, False	46
Everlasting	22	Gooseberry	14	Hemerocallis	44
		Goosefoot	34	Hemlock	17, 41
Fagopyrum	35	Goosefoot Family	34	Ground	41
Fagus	39	Gramineæ	51	Poison	17
Fennel	17	Grape	9	Water	17
Fern, Adder's-tongue	58	Grass, Barnyard	52	Hemp	37
Beech	57	Beard	52	Indian	27
Bladder	57	Bent	53	Henbane	30
Christmas	57	Blue-eyed	44	Hepatica	1
Cinnamon	57	Blue-joint	53	Heracleum	16
Flowering	57	Bottle-brush	55	Herb-Robert	8
Grape	58	Brome	54	Hesperis	4
Lady	58	Bur	52	Hibiscus	7
Maiden-hair	56	Canary	52	Hickory	38
Queen	57	Cat's-tail	53	Hieracium	24
Royal	57	Cotton	49	Hierochloe	52
Sensitive	57	Couch	55	Hippuris	15
Shield	57	Crab	51	Hobble-bush	19
Sweet	38	Darnel	55	Hog Pea-nut	11
Fern Family	56	Drop-seed	53	Holcus	53
Festuca	54	Eel	42	Holly	8
Fever-wort	19	Fescue	54	Mountain	8
Ficoideæ	16	Fox-tail	52, 53	Holly Family	8
Figwort	30	Hair	53, 54	Hone-wort	17
Figwort Family	30	Holy	52	Honeysuckle	19
Filices	56	Indian	52	Honeysuckle Family	19
Fimbrystilis	49	June	54	Hop-tree	8
Fir	41	Mauna	54	Hop-vine	37
Fireweed	15, 23	Meadow	53, 54	Hordeum	55
Five-finger	13	Millet	53	Horehound	33
Flax Family	8	Mountain-rice	52	Hornbeam	38
Fleabane	21	Oat	53, 54	Hornwort Family	40
Floating-heart	28	Old-witch	52	Horse-chestnut	9
Feniculum	17	Orchard	54	Horse-tail Family	56
Forget-me-not	29	Panic	51, 52	Horse-mint	32
Foxglove	31	Poverty	52	Horse-radish	4
Fragaria	12	Reed-bent	53	Horseweed	21
Fraxinus	27	Ribbon	52	Hound's-tongue	28
Fritillaria	46	Rush	53	Houstonia	19
Fringe-tree	27	Rye	55	Huckleberry	25
Frog's-bit Family	42	Spear	54	Humulus	37
Fumaria	3	Sweet vernal	52	Hyacinthus	46
Fumariaceæ	3	Tape	42	Hydrocaridaceæ	42
Fumitory	3	Thin	53	Hydrocotyle	17
Climbing	3	Timothy	53	Hydrophyllæ	28
Fumitory Family	3	Velvet	53	Hydrophyllum	28
Funkia	46	White	52, 54	Hyoscyamus	30
		Wire	54	Hypericaceæ	7
Galeopsis	33	Wood-reed	53	Hypericum	7
Galingale	48	Grass Family	51	Hysop	33
Galmsoga	23	Gratiola	30	Hysopus	33
Galium	20	Greenbrier	44	Iberis	5
Garlic	44	Gromwell	29	Ilex	8
Gapltheria	26	Ground Ivy	33	Ilicineæ	8
Gaylussacia	25	Ground-nut	11	Illecebraceæ	34
Gentiana	28	Groundsel	23	Ilysanthes	30
Gentianaceæ	28			Impatiens	8
Gentian Family	28	Habenaria	43	Indian-corn	55
Geraniaceæ	8	Hackberry	37	Indian-pipe	27
Geranium	8	Haloragieæ	15	India-wheat	35
Geranium Family	8	Hamamelideæ	15	Inula	22
Gerardia	31	Hamamelis	15	Ipomea	29
Germander	32	Hardhack	12	Iridaceæ	44
Geum	12	Harebell	25	Iris	44
Ginseng	17	Hawkweed	24	Iris Family	44
Ginseng Family	17	Hawthorn	13	Ironwood	38
Gladidius	44	Hazelnut	38	Isoetes	58
Gleditschia	12	Hazel, Witch	15	Ivy, Poison	10
Glyceria	54	Heath Family	25	Jacob's ladder	28
Gnaphalium	22	Hedeoma	32	Jamestown-weed	30
Golden-rod	20	Hedge-hyssop	30	Jewel-weed	8
Gold-thread	2	Helianthemum	5		

Juglandaceæ	37	Loosestrife Family	15	<i>Morus</i>	37
Juglans	37	Lopseed	32	Motherwort	33
Juncus	46	Loranthaceæ	36	Mountain-fringe	3
Juncaceæ	46	Lousewort	31	<i>Muhlenbergia</i>	53
June-berry	14	Lovage	17	<i>Mulberry</i>	12, 37
Juniper	41	Lucerne	11	(<i>Mulgedium</i>)	24
Juniperus	41	Ludwigia	15	Mullein	30
		Lungwort	29	<i>Muscari</i>	46
Kalmia	26	Lupinus	10	Mustard	4
Knawel	34	Luzula	46	Mustard Family	4
Knotwort Family	34	Lycium	30	<i>Mycoystylis</i>	42
Krigia	24	Lychnis	6	<i>Myosotis</i>	29
Labiatae	32	<i>Lycopersicum</i>	30	<i>Myrica</i>	38
Lactuca	24	Lycopodium	58	<i>Myriophyllum</i>	15
Ladies'-tresses	42	Lycopus	32		
Lady's-slipper	44	Lysimachia	27	(<i>Nabalus</i>)	24
Lady's-thumb	35	Lythraceæ	15	<i>Naiadaceæ</i>	48
Lagenaria	16	Lythrum	15	<i>Najas</i>	48
Lamium	33			(<i>Nardosmia</i>)	23
Laportea	37	Madder Family	19	<i>Nasturtium</i>	4
(Lappa)	23	Maianthemum	45	<i>Negundo</i>	9
Larch	41	Maize	55	<i>Nemopanthes</i>	8
Larix	41	Mallow Family	7	<i>Nepea</i>	33
Larkspur	2	Malva	1	Nettle	33, 37
Lauraceæ	36	Malvaceæ	7	Nettle Family	37
Laurel	26	Mandrake	3	<i>Nicandra</i>	30
Laurel Family	36	Maple	9	<i>Nicotiana</i>	30
Larandula	33	Maple, Ash-leaved	9	Nightshade, Enchanters'	16
Lavender	33	Mare's tail	15	Nightshade Family	29
Leather-leaf	26	Marigold, Bur	22	<i>Nuphar</i>	3
Leatherwood	36	Marsh	2	<i>Nymphaea</i>	3
Lechea	5	Water	23	<i>Nymphaeaceæ</i>	3
Ledum	26	<i>Marjoram</i>	32	<i>Nyssa</i>	18
Leek, Wild	44	Marrubium	33		
Leersia	52	Matrimony-vine	30	Oak	38
Leguminosæ	10	<i>Mothiola</i>	5	<i>Jerusalem</i>	34
Lemna	47	May-flower	26	<i>Oakesia</i>	45
Lemmaceæ	47	May-weed	23	Oak Family	38
Lentibulaceæ	31	Meadow-sweet	12	<i>Ocimum</i>	33
Leonurus	33	Medeola	45	<i>Oenothera</i>	16
Lepidium	5	Medicago	11	Oleaceæ	27
Lespedeza	11	Medick	11	Olive Family	27
Lettuce	24	Melampyrum	31	<i>Onagraceæ</i>	15
Leverwood	38	Melilotus	10	Onion	44
Ligusticum	17	<i>Melissa</i>	33	<i>Onoclea</i>	57
Ligustrum	27	Menispermum	2	<i>Ophioglossaceæ</i>	58
Lilac	27	Menispermaceæ	2	<i>Ophioglossum</i>	58
Liliaceæ	44	Mentha	32	Orchidaceæ	42
Lilium	45	Menyanthes	28	Orchis	43
Lily	3, 44, 45, 46	Mercury, Three-seeded	36	<i>Orchis</i> Family	42
Day	44, 46	<i>Mertensia</i>	29	<i>Origanum</i>	32
Sword	44	Mermaid-weed	15	<i>Ornithogalum</i>	44
Tiger	45	<i>Mezereum</i> Family	36	<i>Orobanchaceæ</i>	31
Water	3	Milkweed Family	27	<i>Orpine</i> Family	15
Lily Family	44	Milkwort Family	10	<i>Oryzopsis</i>	52
Lily of the Valley	44	Milium	53	<i>Ostrya</i>	38
Limnanthemum	28	Mimulus	30	<i>Osmorrhiza</i>	17
Linaceæ	8	Mint Family	32	<i>Osmunda</i>	57
Linaria	30	Mistletoe Family	36	Oyster-plant	24
Linden Family	7	Mitella	14	<i>Oxalis</i>	8
Lindera	36	Mitchella	20	Ox-eye	22
Linnaea	19	Mitre-wort	14		
Linum	8	Mock-orange	14	<i>Paeonia</i>	2
Liparis	42	Mollugo	16	<i>Panicum</i>	51
Listera	42	Monarda	32	<i>Papaveraceæ</i>	3
Lithospermum	29	Moneses	26	<i>Parnassia</i>	14
Liverwort	1	Moneywort	27	<i>Parsety</i>	17
Lobeliaceæ	25	Monkey-flower	30	<i>Parsley</i> Family	16
Lobelia Family	25	Monkshood	2	Parsnip	16, 17
Logust	11, 12	<i>Monotropa</i>	27	Partridge-berry	20
Loiseleuria	26	Moonseed	2	<i>Paspalum</i>	51
Lolium	55	Moonseed Family	2	<i>Pastinaca</i>	16
Lonicera	19	Moonwort	58	<i>Pea</i>	12
Loosestrife	15, 27	Morning-glory	29	<i>Pea-nut</i>	12

<i>Pear</i>	14	<i>Potato</i>	30	<i>Salix</i>	39
<i>Pearlwort</i>	6	<i>Potentilla</i>	13	<i>Salvia</i>	33
<i>Pea-tree</i>	12	<i>Poterium</i>	13	<i>Sambucus</i>	19
<i>Pedicularis</i>	31	<i>Prenanthes</i>	24	<i>Sandalwood Family</i>	36
<i>Pellæa</i>	56	<i>Primrose, Evening</i>	16	<i>Sandwort</i>	6
<i>Peltandra</i>	47	<i>Primrose Family</i>	27	<i>Sanguinaria</i>	3
<i>Pennyroyal</i>	32	<i>Primulaceæ</i>	27	<i>Sanicula</i>	17
<i>Pennywort, Water</i>	17	<i>Prince's-feather</i>	35	<i>Santalaceæ</i>	36
<i>Penthorum</i>	15	<i>Prince's pine</i>	26	<i>Sapindaceæ</i>	9
<i>Pentstemon</i>	30	<i>Privet</i>	27	<i>Saponaria</i>	6
<i>Peppergrass</i>	5	<i>Proserpinaca</i>	15	<i>Sarracenia</i>	3
<i>Pepperidge</i>	18	<i>Prunus</i>	12, 14	<i>Sarraceniaceæ</i>	3
<i>Peppermint</i>	32	<i>Ptelea</i>	8	<i>Sarsaparilla</i>	17
<i>Pepper-root</i>	4	<i>Pteris</i>	56	<i>Sassafras</i>	36
<i>Petasites</i>	23	<i>Pterospora</i>	26	<i>Satureia</i>	33
<i>Phalaris</i>	52	<i>Pulse Family</i>	10	<i>Saxifraga</i>	14
<i>Phaseolus</i>	12	<i>Purslane</i>	7	<i>Saxifragaceæ</i>	14
<i>Phegopteris</i>	57	<i>Purslane Family</i>	7	<i>Saxifrage Family</i>	14
<i>Philadelphus</i>	14	<i>Putty-root</i>	42	<i>Scheuchzeria</i>	48
<i>Phleum</i>	53	<i>Pycnanthemum</i>	32	<i>Scilla</i>	46
<i>Phloe</i>	28	<i>Pyrola</i>	26	<i>Scirpus</i>	49
<i>Phryma</i>	32	<i>Pyrus</i>	13, 14	<i>Scleranthus</i>	34
<i>Physalis</i>	29			<i>Serophularia</i>	30
<i>Physostegia</i>	33			<i>Serophulariaceæ</i>	30
<i>Phytolacca</i>	35			<i>Scutellaria</i>	33
<i>Phytolaccaceæ</i>	35			<i>Scutellaria</i>	55
<i>Picea</i>	41			<i>Sedge</i>	49
<i>Pickerel-weed Family</i>	46			<i>Sedge Family</i>	48
<i>Pig-nut</i>	38			<i>Sedum</i>	15
<i>Pig-weed</i>	34			<i>Selaginella</i>	58
<i>Pilea</i>	37			<i>Selaginellaceæ</i>	58
<i>Pimpernel</i>	27, 30	<i>Ranunculus</i>	2	<i>Selaginella Family</i>	58
<i>Pine</i>	41, 58	<i>Raphanus</i>	5	<i>Selt-heal</i>	33
<i>Pine-drops</i>	26	<i>Raspberry</i>	12	<i>Senecio</i>	23
<i>Pine Family</i>	41	<i>Rattlesnake-plantain</i>	43	<i>Senna</i>	11
<i>Pine sap</i>	27	<i>Rattlesnake-weed</i>	24	<i>Sericocarpus</i>	21
<i>Pink Family</i>	6			<i>Setaria</i>	52
<i>Pinus</i>	41			<i>Shad-bush</i>	14
<i>Pinweed</i>	5			<i>Sheep-berry</i>	19
<i>Pipe-vine</i>	36	<i>Rhododendron</i>	26	<i>Shepherd's-purse</i>	4
<i>Pipewort Family</i>	48	(<i>Rhodora</i>)	26	<i>Sicyos</i>	16
<i>Pisum</i>	12	<i>Rhubarb</i>	35	<i>Silene</i>	6
<i>Pitcher-plant</i>	3	<i>Rhus</i>	10	<i>Sisymbrium</i>	4
<i>Pitcher-Plant Family</i>	3	<i>Rhynchospora</i>	49	<i>Sisyrinchium</i>	44
<i>Plane-tree Family</i>	37	<i>Ribes</i>	14	<i>Sium</i>	17
<i>Plantaginaceæ</i>	33	<i>Ricinus</i>	37	<i>Skullcap</i>	33
<i>Plantago</i>	33	<i>Robinia</i>	11	<i>Smartweed</i>	35
<i>Plantain Family</i>	33	<i>Rocket</i>	4	<i>Smilacina</i>	45
<i>Platanaceæ</i>	37	<i>Rock-rose</i>	5	<i>Suirax</i>	44
<i>Platanus</i>	37	<i>Rock-rose Family</i>	5	<i>Snake-head</i>	30
<i>Plum</i>	12	<i>Rosa</i>	13	<i>Snakeroot</i>	17, 20
<i>Poa</i>	54	<i>Rosaceæ</i>	12	<i>Snowberry</i>	19
<i>Podophyllum</i>	3	<i>Rose Family</i>	12	<i>Creeping</i>	25
<i>Pogonia</i>	43		34	<i>Soapwort</i>	6
<i>Poke, Indian</i>	46		19	<i>Soapberry Family</i>	9
<i>Pokeweed Family</i>	35	<i>Rubiaceæ</i>	12	<i>Solanaceæ</i>	29
<i>Polypodium</i>	56	<i>Rudbeckia</i>	22	<i>Solanum</i>	20, 30
<i>Polemoniaceæ</i>	28	<i>Rue, Meadow</i>	1	<i>Solidago</i>	29
<i>Polemonium</i>	28		1	<i>Solomon's-seal</i>	44, 45
<i>Polemonium Family</i>	28	<i>Rumex</i>	35	<i>Sonchus</i>	24
<i>Polygala Family</i>	16	<i>Rush Family</i>	46	(<i>Sorghum</i>)	52
<i>Polygalaceæ</i>	10	<i>Rush</i>	46	<i>Sorrel</i>	8, 35
<i>Polygonaceæ</i>	35	<i>Beak</i>	49	<i>Sour-gum tree</i>	18
<i>Polygonatum</i>	44	<i>Club</i>	49	<i>Sparganium</i>	47
<i>Polygonum</i>	35	<i>Spike</i>	49	<i>Spearmint</i>	32
<i>Pondweed</i>	48	<i>Twig</i>	49	<i>Specularia</i>	25
<i>Pondweed Family</i>	48	<i>Rutaceæ</i>	8	<i>Speedwell</i>	31
<i>Pontederia</i>	46	<i>Rye</i>	55	<i>Spergula</i>	7
<i>Pontederiaceæ</i>	46	<i>Wild</i>	55	(<i>Spergularia</i>)	6
<i>Poplar</i>	39			<i>Spice-wood</i>	36
<i>Poppy Family</i>	3	<i>Sage</i>	33	<i>Spikenard</i>	17
<i>Populus</i>	39	<i>Sagina</i>	6	<i>Spinacia</i>	34
<i>Portulaca</i>	7	<i>Sagittaria</i>	47	<i>Spiraea</i>	12, 14
<i>Portulacaceæ</i>	7		7	<i>Spiranthes</i>	42
<i>Potamogeton</i>	48	<i>St. John's-wort Family</i>	7	<i>Spirodela</i>	47
		<i>Salicaceæ</i>	39		

Spleenwort	56	Thoroughwort	20	Veronica	31
Sporobolus	53	Thuya	41	Vervain Family	32
Spring-beauty	7	Thyme	33	Vetch	11
Spruce	41	Thymelæcæ	36	Viburnum	19
Spurge Family	36	Thymus	33	Vicia	11
Spurrey	6, 7	Tiarella	14	Vincetoxicum	28
Squirrel-corn	3	Tiger-flower	44	Vine Family	9
Stachys	33	Tigridia	44	Viola	5
Staff-tree Family	8	Tilia	7	Violaceæ	5
Staphylea	10	Tiliaceæ	7	Violet Family	5
Star-flower	27	Toad-Flax	30, 36	Virginian creeper	9
Star-of-Bethlehem	44	Tobacco	30	Virgin's-bower	1
Starwort	21	Tobacco, Indian	25	Vitaceæ	9
Water	15	Totifeldia	46	Vitis	9
Steironema	27	Tomato	30		
Stellaria	6	Tragopogon	24	Wake-robin	45
Stock	5	Trefoil	19	Walking-leaf	57
Stone-crop	15	Tick	11	Waldsteinia	12
Storks-bill	8	Trichostema	32	Wainut Family	37
Stramonium	30	Trientalis	27	Waterleaf Family	28
Strawberry	12	Trifolium	10	Water-lily Family	3
Streptopus	45	Trillium	45	Water-milfoil Family	15
Sundew	15	Triosteum	19	Water-plautain Family	47
Sundew Family	15	Triticum	55	Water-shield	3
Sunflower	22	Tsuga	41	Waterweed	42
Sumach	10	Tulipa	46	Wheat	55
Summer Savory	33	Tumbleweed	34	Cow	31
Sweet- Cicely	17	Turnip	5	Wild Ginger	36
Sweet-fern	38	Indian	47	Willow	39
Sweet-flag	47	Tussilago	23	Willow Family	39
Sweet-gale	38	Twayblade	42	Willow-herb	15
Sweet-gale Family	38	Twin-flower	19	Wind-flower	1
Sycamore	37	Twist-foot	45	Wintergreen	26
Symporicarpus	19	Typha	47	Wistaria	12
Sympythium	29	Typhaceæ	47	Witch-hazel Family	15
Symplocarpus	47			Woodbine	9
Syringa	14, 27	Ulmus	37	Woodisia	57
		Umbelliferæ	16	Wormwood	23
Tamarack	41	Urtica	37		
Tanacetum	23	Urticaceæ	37	Xanthium	22
Tansy	23	Utricularia	31	Xanthoxylum	8
Taraxacum	24	Uvularia	45	Xyridaceæ	46
Taxus	41			Nyris	46
Tea, Labrador	26	Vaccinium	25	Yarrow	23
New Jersey	9	Valerian, Greek	28	Yellow-eyed-grass Family	46
Teucrium	32	Vallisneria	42	Yew	41
Thalictrum	1	Venus's looking-glass	25	Yucca	46
(Thaspium)	17	Veratrum	46		
Thistle	24, 29	Verbascum	30	Zannichellia	48
Thlaspi	4	Verbena	32	Zea	55
Thorn-tree	13	Verbenaceæ	32	Zizia	17


INDEX TO VERTEBRATES.

Acanthis	72	Butcher-bird	74	Crotalus	63
Acanthopteri	60			Crow	71
Accipiter	68	Calcarius	72	Cuckoo	69
Acris	61	Calomys	78	Cuculidæ	69
Actitis	67	Canidae	80	Cyanocitta	71
Adder	63	Caprimulgidae	70	Cyprinidæ	59
Ægialitis	67	Cariacus	80		
Agelaius	72	Carnivora	80	Dace	59
Aix	66	Carpodacus	72	Darters	60
Alaudidae	71	Castor	78	Deer	80
Alcedinidæ	69	Castoridae	78	Dendragapus	68
Alcidæ	65	Cat-bird,	76	Dendroica	75
Alle	65	Cat-fish.	59	Diadophis	63
Amblystoma	61	Catostomidae	59	Diemyctylus	61
Amblystomatidæ	61	Catostomus	59	Dipper-duck	65
Ameiurus	59	Cedar-bird,	74	Diving-birds	65
Ammodramus	72, 73	Centrarchidae	60	Dolichonyx	71
Ampelidæ	74	Ceophleens	70	Dove	68
Ampelis	74	Certbia	76	Dryobates	70
Anas	66	Certhiidæ	76	Duck	66
Anatidæ	66	Cervidae	80		
Anguilla	60	Ceryle	69	Eagle	68
Anguillidæ	60	Chætura	70	Ectopistes	68
Anseres	66	Charadriidæ	67	Eel	60
Anthus	75	Chelidon	73	Eel Pout	60
Antrostomus	70	Cheloporus	64	Empidonax	71
Apodes	60	Chelydra	63	Emydidae	64
Aquila	68	Chelydridæ	63	Erethizon	78
Archibuteo	68	Chewink	73	Erismatura	66
Arctomyys	79	Chickadee	76	Esocidae	60
Ardea	66	Chimney Swallow	70	Esox	60
Ardeidæ	66	Chipmunk	79	Etheostoma	60
Arenaria	67	Chiroptera	79	Eutainia	63
Arvicola	78	Chordeiles	70	Eventognathi	59
Asio	69	Chorophylus	62	Evet	61
Atalapha	79	Chrysomys	64	Evotomys	78
Auk	65	Chub Eel	60		
Aves	65	Circus	68	Falco	68, 69
		Cistothorus	76	Falcon	68
Bartramia	67	Clivicola	74	Falconidæ	68
Bass	60	Coccyges	69	Felidae	80
Bat	70	Coccyzus	69	Felis	80
Batrachia	61	Colaptes	70	Ferae	80
Bear	80	Colinus	67	Fiber	78
Beaver	78	Colubridæ	63	Finch	72
Bittern	66	Columbae	68	Flycatcher	71
Blackbird	71, 72	Columbidæ	68	Fox	80
Black Cat	80	Colymbus	65	Fringillidæ	72
Blarina	79	Compsothlypis	75	Frog	61, 62
Blue Bird	77	Condylura	79	Tree	62
Bobolink	71	Contopus	71	Fulica	67
Bob-white	67	Cormorant	65		
Bonasa	68	Coot	67	Gadidæ	60
Botaurus	66	Corvidæ	71	Galeoscoptes	76
Branta	66	Corvus	71	Gallinae	67
Brown Thrasher	76	Cottidae	60	Gallinago	67
Bubo	69	Cottus	60	Geothlypis	75
Bubonidæ	69	Cow-bird	71	Glires	78
Bufo	61	Crane	66	Golden Robin	72
Bufo	61	Creeper	74, 76	Goldfinch	72
Bull-head	59	Crossbill	72	Goose	66
Buteo	68	Crotalidae	63	Goshawk	68

Grebe	65	Micropterus	60	Podiceps	65
Grosbeak	72, 73	Miller's Thumb	60	Podicipidae	65
Grouse	67, 68	Mimus	76	Podilymbus	65
Gull	65	Mink	80	Pooecetes	72
Gulo	8)	Minnow	59	Pörenpine	78
		Mniotilla	74	Porzana	67
Habia	73	Mniotillidæ	74	Primates	81
Haliaëtus	68	Mocking-bird	76	Procyon	80
Haplopi	60	Mole	79	Procyonidæ	80
Hare	78	Molothrus	71	Progne	73
Harporhynchus	76	Motacillidæ	75	Pumpkin-seed	60
Hawk	68, 69	Mud Hen	67	Purple-finck	72
Hedgehog	78	Muridæ	78	Putorius	80
Helminthophila	74	Mus	78	Pygopodes	65
Herodiones	66	Muskrat	78		
Heron	66	Mustela	80	Quail	67
Heterodon	63	Mustelidæ	80	Quiscalus	72
High-holer	70	Myarchus	71	Rabbit	78
Hirundinidæ	73			Raccoon	80
Hominidae	81	Nematognathi	59	Rai	66, 67
Homo	81	Newt	61	Rallidæ	66
Horn-pout	59	Night-hawk	70	Rallus	66
Humming-bird	70	Notemigonous	59	Rana	62
Hyla	62	Notropis	59	Ranidæ	62
Hyliidae	61	Nuthatch	76	Raptores	68
Hystricidæ	78	Nyctala	69	Rat	75
		Nyctea	69	Rattlesnake	66
Icteridæ	71	Nycticorax	66	Red-breast	77
Icterus	72		63	Red-poll-linnet	72
Indian Hen	66	Ophibolus	63	Redstart	75
Indi.-o-bird	73	Ophidia	72	Regulus	76
Insectivora	79	Oriole	69	Rerilia	63
Isopondyli	60	Osprey	71	Rhinichthys	59
Jay	71	Otocoris	75	Robin	72, 73, 77
Jumping Mouse	78	Oven-bird	69	Rodents	78
Junco	73	Owl			
Kildeer	67	Paludicolaæ	66	Sable	80
King-bird	71	Pandion	69	Salamander	61
Kingfisher	69	Panther	80	Salientia	61
Kinglet	76	Paride	76	Salmonidæ	60
		Partridge	68	Salmon-trout	60
Laniidae	74	Parus	76	Salvelinus	60
Lanius	74	Passer	72	Sand-martin	74
Lapland-longspur	72	Passeres	71	Sandpiper	67
Laridæ	65	Passerella	73	Sap-sucker	70
Lark	71, 75	Passerina	73	Sayornis	71
Larus	65	Pelobatidæ	61	Scalops	73
Lepomis	60	Perca	60	Scapanus	79
Leporidæ	78	Perch	60	Scaphiopus	61
Lepus	78	Percidæ	60	Seiruridae	79
Limicolæ	67	Perisoreus	71	Sciuropterurus	79
Liopeltis	63	Petrochelidon	73	Seirus	79
Longipennes	65	Pewee	71	Scolecopagrus	72
Loon	65	Phalaenocoracidae	65	Scolopacidae	67
Lophodytes	66	Phalaenocorax	65	Scotiaptex	69
Lota	60	Philohela	67	Sculpin	60
Loxia	72	Phoebe	71	Seirus	75
Lynx	80	Pieciæ	70	Semotilus	59
		Pickerel	70	Setophaga	75
Macrochires	70	Picoides	60	Sheldrake	66
Mammalia	78	Pigeon, Wild	70	Shiner	59
Man	81	Pike	68	Shrew	79
Marsh-Harrier	68	Pine-finch	60	Shrike	74
Martin, Purple	73	Pine-marten	72	Sialia	77
Meadow-lark	72	Pinicola	80	Siluridae	59
Megascops	69	Pipilo	72	Sitta	76
Melanerpes	70	Piranga	73	Skunk	80
Melospiza	78	Pisces	73	Snake	63
Mephitis	80	Plectrophenax	59	Snipe	67
Merganser	66	Plethodon	72	Snow-bird	73
Merula	77	Plethodontidæ	61	Snow-bunting	72
Mice	78,	Pleurodelidæ	61	Sorex	79
Micropodidæ	70	Plover	61	Sorcidæ	79
			67	Spade-foot	61

INDEX.

Sparrow	72, 73	Testudinata	63	Ursus	80
Spelerves	61	Tetraonidæ	67		
Sphyrapicus	70	Thrush	75, 76, 77	Vespertilio	79
Spinus	72	Tip up	67	Vespertilionidæ	79
Spizella	73	Titlark	75	Vesperugo	79
Squirrel	79	Titmouse	76	Vireo	74
Squawk	66	Toad	61, 62	Vireonidæ	74
Sterna	65	Tree	62	Vulpes	80
Stizostedion	60	Tortoise	64		
Storeria	63	Totanus	67	Wall-eye	60
Sturnella	72	Tringa	67	Warbler	74, 75, 76
Sucker	59	Trochilidæ	70	Water-wagtail	75
Sunfish	60	Trochilus	70	Wax-wing	74
Surnia	69	Troglodytes	76	Weasel	80
Swallow	73, 74	Troglodytidæ	76	Whippoorwill	70
Swift	70	Tropidonotus	63	Wild-cat	80
Sylvania	75	Trotut	60	Wolverene	80
Sylviidæ	76	Turdidæ	76	Woodchuck	79
Syrnium	69	Turdus	76, 77	Woodcock	67
		Turnstone	67	Woodpecker	70
Tachycineta	73	Turtle	63, 64	Wren	76
Talpidae	79	Tyranidæ	71		
Tamias	79	Tyrannus	71	Yellow-bird	72
Tanager	73			Yellow-hammer	70
Tanagridæ	73	Ungulata	80	Yellow-throat	75
Tattler	67	Urinator	65	Zapodidæ	78
Teal	66	Urinatoridæ	65	Zapus	78
Tell-tale	67	Urodela	61	Zenaidura	68
Tern	65	Ursidæ	80	Zonotrichia	73



A223

